

Novel imidazolinone derivatives of formula (I), which are useful as angiotensin II antagonists, are disclosed.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MN	Mongolia
AU	Australia	FR	France	MR	Mauritania
BB	Barbados	GA	Gabon	MW	Malawi
BE	Belgium	GB	United Kingdom	NL	Netherlands
BF	Burkina Faso	GN	Guinea	NO	Norway
BG	Bulgaria	GR	Greece	NZ	New Zealand
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	PT	Portugal
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	RU	Russian Federation
CG	Congo	KP	Democratic People's Republic of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SE	Sweden
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovak Republic
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CS	Czechoslovakia	LU	Luxembourg	SU	Soviet Union
CZ	Czech Republic	MC	Monaco	TD	Chad
DE	Germany	MG	Madagascar	TG	Togo
DK	Denmark	MI	Mali	UA	Ukraine
ES	Spain			US	United States of America

TITLE

ANGIOTENSIN II RECEPTOR
BLOCKING IMIDAZOLINONE DERIVATIVES

CROSS-REFERENCE TO RELATED APPLICATION

This application is a continuation-in-part of U.S.
Application Serial Number 07/747,023, filed August 19,
1991.

BACKGROUND OF THE INVENTIONField of the Invention

15

This invention relates to novel substituted
imidazolinone derivatives. The invention also relates
to pharmaceutical compositions containing the novel
imidazolinone derivatives and pharmaceutical methods
using them, alone and in conjugation with other drugs.

The compounds of this invention inhibit the action
of the hormone angiotensin II (AII) and are useful
therefore in alleviating angiotensin induced
hypertension. The enzyme renin acts on a blood plasma
 α 2-globulin, angiotensinogen, to produce angiotensin I,
which is then converted by ACE to AII. The latter
substance is a powerful vasopressor agent which has been
implicated as a causative agent for producing high blood
pressure in various mammalian species, such as the rat,
dog, and man. The compounds of this invention inhibit
the action of AII at its receptors on target cells and
thus prevent the increase in blood pressure produced by
this hormone-receptor interaction. By administering a
compound of this invention to a species of mammal with


hypertension due to AII, the blood pressure is reduced. The compounds of this invention are also useful for the treatment of congestive heart failure. Administration of a compound of this invention with a diuretic such as furosemide or hydrochlorothiazide, either as a stepwise combined therapy (diuretic first) or as a physical mixture, enhances the antihypertensive effect of the compound. Administration of a compound of this invention with a NSAID can prevent renal failure which sometimes results from administration of a NSAID.

Several peptide analogs of AII are known to inhibit the effects of this hormone by competitively blocking the receptors, but their experimental and clinical applications have been limited by the partial agonist activity and lack of oral absorption (M. Antonaccio, *Clin. Exp. Hypertens.*, **1982**, A4, 27-46; D. H. P. Streeten and G. H. Anderson, Jr., Handbook of Hypertension, Clinical Pharmacology of Antihypertensive Drugs, ed., A. E. Doyle, Vol. 5, pages 246-271, Elsevier Science Publisher, Amsterdam, The Netherlands, 1984).

Several non-peptide antagonists of AII have been disclosed. These compounds are covered by U.S. Patents 4,207,324; 4,340,598; 4,576,958; 4,582,847; and 4,880,804; in European Patent Applications 028,834; 245,637; 253,310; and 291,969; and in articles by A. T. Chiu, et al. (*Eur. J. Pharm. Exp. Therap.*, **1988**, 157, 13-21) and by P. C. Wong, et al. (*J. Pharm. Exp. Therap.*, **1988**, 247, 1-7). All of the U.S. Patents, European Patent Applications 028,834 and 253,310 and the two articles disclose substituted imidazole compounds which are generally bonded through a lower alkyl bridge to a substituted phenyl. European Patent Application 245,637 discloses derivatives of 4,5,6,7-tetrahydro-2H-imidazo[4,5-c]pyridine-6-carboxylic acid and analogs


thereof as antihypertensive agents, specifically Ca^{2+} channel blockers.

L. Chang et al., in EP 0 412 594 A (filed July 23, 1990) disclose substituted triazolinones, triazolinethiones, and triazolinimines of the formula:




These are claimed to be antagonists of AII which are useful for treating hypertension, congestive heart failure (CHF), and elevated intraocular pressure.

C. Bernhart et al., in WO 91/14679 (published October 3, 1991) disclose heterocyclic N-substituted derivatives of the formula




These compounds are disclosed to be antagonists of AII which are useful for treating cardiovascular disorders such as hypertension.

F. Ostermeyer et al., in EP 475,898 (published March 18, 1992) disclose heterocyclic N-substituted derivatives of formula



These compounds are disclosed to be antagonists of AII which are useful for treating cardiovascular disorders such as hypertension.

P. Herold and P. Bühlmyer in EP 0 407 342 A2 disclose substituted pyrimidinones, pyrimidinethiones, and pyrimidinimines of the formula:




These are claimed to be antagonists of AII which are useful for treating hypertension.

E. Allen, et al. in EP 0 419 048 A (filed August 21, 1990) disclose a similar series of pyrimidinones which are claimed to be antagonists of AII

useful for the treatment of CHF and elevated intraocular pressure.

SUMMARY OF THE INVENTION


5 The present invention provides novel angiotensin II receptor antagonists of formula (I), pharmaceutical compositions containing compounds of formula (I) and therapeutic methods using them



10

wherein:

R¹ is other than in the ortho position and is:

15 R² is

- (a) H,
- (b) halo (F, Cl, Br, I),
- (c) C₁-C₄ alkyl,
- (d) C₁-C₄ alkoxy,
- 20 (e) C₁-C₄ acyloxy,
- (f) C₁-C₄ alkylthio,
- (g) C₁-C₄ alkylsulfinyl,
- (h) C₁-C₄ alkylsulfonyl,
- (i) hydroxy (C₁-C₄) alkyl,
- 25 (j) aryl (C₁-C₄) alkyl,

- (k) -CO₂H,
(l) -CN,
(m) -CONHOR¹²,
(n) -SO₂NHR²¹,
5 (o) -NH₂,
(p) C₁-C₄ alkylamino,
(q) C₁-C₄ dialkylamino,
(r) -NHSO₂R²⁰,
(s) -NO₂,
10 (t) furyl,
(u) aryl;

wherein aryl is phenyl optionally substituted with one or two substituents selected from the group consisting of halo, C₁-C₄ alkyl, C₁-C₄ alkoxy, -NO₂,
15 -CF₃, C₁-C₄ alkylthio, -OH, -NH₂, C₁-C₄ alkylamino, C₁-C₄ dialkylamino, -CN, -CO₂H, -CO₂CH₃, -CO₂CH₂CH₃, -CO₂-benzyl;

R³ is

- (a) H,
20 (b) halo,
(c) C₁-C₄ alkyl,
(d) C₁-C₄ alkoxy,
(e) C₁-C₄ alkoxyalkyl;

R⁴ is

- 25 (a) -CN,
(b) -NO₂,
(c) -CO₂R¹¹;

R⁵ is

- (a) H,
30 (b) C₁-C₆ alkyl,
(c) C₃-C₆ cycloalkyl,
(d) C₂-C₄ alkenyl,
(e) C₂-C₄ alkynyl;

R⁶ is

- (a) C₁-C₁₀ alkyl,
- (b) C₃-C₈ alkenyl,
- (c) C₃-C₈ alkynyl,
- 5 (d) C₃-C₈ cycloalkyl,
- (e) C₄-C₈ cycloalkenyl,
- (f) C₄-C₁₀ cycloalkylalkyl,
- (g) C₅-C₁₀ cycloalkylalkenyl,
- (h) C₅-C₁₀ cycloalkylalkynyl,
- 10 (i) $-(CH_2)_5Z(CH_2)_mR^5$,
- (j) phenyl, optionally substituted with 1-2 substituents selected from the group of halo, C₁-C₄ alkyl, C₁-C₄ alkoxy, -NO₂, -NH₂, -OH and benzyloxy,
- (k) benzyl, optionally substituted on the phenyl
- 15 ring with 1-2 substituents selected from the group of halo, C₁-C₄ alkyl, C₁-C₄ alkoxy and -NO₂;

R⁷, R⁸, R⁹, and R¹⁰ are independently chosen from

- (a) H,
- 20 (b) C₁-C₈ alkyl,
- (c) C₁-C₈ perfluoroalkyl,
- (d) C₃-C₆ cycloalkyl,
- (e) -NO₂,
- (f) -CN,
- 25 (g) -CONR¹⁵R¹⁶,
- (h) -CO₂R¹⁷,
- (i) -OR¹⁸,
- (j) $-(CH_2)_nCONR^{15}R^{16}$ where n is 1-4,
- (k) $-(CH_2)_nCO_2R^{17}$ where n is 1-4,
- 30 (l) $(CH_2)_nOR^{18}$ where n is 1-4,
- (m) aryl, wherein aryl is as defined above,
- (n) -CH₂ aryl, wherein aryl is as defined above,
- (o) R⁷ and R⁸ taken together are $-(CH_2)_t-$, or $-(CH_2)_mX(CH_2)_q-$,

(p) R^9 and R^{10} taken together can be S, or O;

R^{11} is


- (a) H,
 (b) C_1-C_4 alkyl,
 5 (c) C_1-C_4 cycloalkyl,
 (d) phenyl,
 (e) benzyl;

R^{12} is


- (a) H,
 10 (b) methyl,
 (c) benzyl;

R^{13} is


- (a) $-CH_2CO_2H$,
 (b) $-C(CF_3)_2OH$,
 15 (c) $-CONHNHSO_2CF_3$,
 (d) $-CONHOR^{12}$,
 (e) $-CONHSO_2R^{20}$,
 (f) $-CONHSO_2NHR^{19}$,
 (g) $-C(OH)R^{19}PO_3H_2$,
 20 (h) $-NHCONHSO_2R^{20}$,
 (i) $-NHPO_3H_2$,
 (j) $-SO_2NHCOR^{19}$,
 (k) $-OPO_3H_2$,
 (l) $-OSO_3H$,
 25 (m) $-PO(OH)R^{19}$,
 (n) $-PO_3H_2$,
 (o) $-SO_3H$,
 (p) $-SO_2NHR^{19}$,
 (q) $-NHSO_2NHCOR^{19}$,
 30 (r) $-SO_2NHCONHR^{19}$,
 (s)



(t)




(u)



5

(v)

R¹⁴ is

- (a) H,
- (b) C₁-C₄ alkyl,
- 10 (c) aryl, wherein aryl is defined as above,
- (d) benzyl,
- (e) COR¹¹,
- (f) CONHR¹¹;

R¹⁵ and R¹⁶ are independently

- 15 (a) H,
- (b) C₁-C₆ alkyl,
- (c) aryl, wherein aryl is as defined above,
- (d) aryl (C₁-C₄) alkyl, where aryl is as defined above,

20 or taken together constitute a

- (e) piperidine ring,
- (f) morpholine ring,
- (g) piperazine ring, optionally N-substituted with C₁-C₆ alkyl, phenyl or benzyl;

25 R¹⁷ is

- (a) H,
- (b) C₁-C₆ alkyl,
- (c) phenyl,
- (d) benzyl;

R¹⁸ is

- (a) H,
- (b) C₁-C₆ alkyl,
- (c) phenyl,
- 5 (d) benzyl;

R¹⁹ is

- (a) H,
 - (b) C₁-C₅ alkyl optionally substituted with a
substituent selected from the group consisting of aryl,
10 where aryl is as defined above, heteroaryl where
heteroaryl is as defined above, -OH, -SH, C₁-C₄ alkyl,
C₁-C₄ alkoxy, C₁-C₄ alkylthio, -CF₃, halo, -NO₂, -CO₂H,
-CO₂CH₃, -CO₂-benzyl, -NH₂, C₁-C₄ alkylamino, C₁-C₄
dialkylamino, -PO₃H₂,
 - 15 (c) aryl, where aryl is defined as above,
 - (d) -CH₂ aryl, where aryl is defined as above,
 - (e) heteroaryl,
- wherein heteroaryl is an unsubstituted,
monosubstituted or disubstituted 5- or 6-membered
20 aromatic ring which can optionally contain from 1 to 3
heteroatoms selected from the group consisting of O, N,
and S and wherein the substituents are members selected
from the group consisting of -OH, -SH, C₁-C₄ alkyl, C₁-C₄
alkoxy, -CF₃, halo, -NO₂, -CO₂H, -CO₂CH₃, -CO₂-benzyl,
25 -NH₂, C₁-C₄ alkylamino, C₁-C₄ dialkylamino;

R²⁰ is

- (a) aryl, where aryl is defined as above,
- (b) C₃-C₇ cycloalkyl,
- (c) C₁-C₄ perfluoroalkyl,
- 30 (d) C₁-C₄ alkyl optionally substituted with a
substituent selected from the group consisting of aryl,
where aryl is as defined above, heteroaryl where
heteroaryl is as defined above, -OH, -SH, C₁-C₄ alkyl,
C₁-C₄ alkoxy, C₁-C₄ alkylthio, -CF₃, halo, -NO₂, -CO₂H,

-CO₂CH₃, -CO₂-benzyl, -NH₂, C₁-C₄ alkylamino, C₁-C₄
dialkylamino, -PO₃H₂,

(e) heteroaryl, where heteroaryl is as defined
above;

5 R²¹ is

- (a) H,
- (b) C₁-C₆ alkyl,
- (c) phenyl,
- (d) benzyl,

10 or taken together constitute a

- (e) piperidine ring,
- (f) morpholine ring,
- (g) piperazine ring, optionally N-substituted with
C₁-C₆ alkyl, phenyl or benzyl;

15

X is

- (a) S,
- (b) O,
- (c) SO,
- 20 (d) SO₂,
- (e) CHR¹⁴,
- (f) NR¹⁴;

Z is

- 25 (a) -O-,
- (b) -S-,
- (c) -NR¹¹-;

m is 1 to 5;

n is 1 to 4;

30 q is 1 to 5;


t is 2 to 5;

or a pharmaceutically acceptable salt thereof.

Preferred compounds of this invention are those of formula (I) wherein

R¹ is in the para position and is

5



R⁶ is

- (a) C₁-C₁₀ alkyl,
- 10 (b) C₃-C₁₀ alkenyl,
- (c) C₃-C₁₀ alkynyl,
- (d) C₃-C₈ cycloalkyl,
- (e) phenyl, optionally substituted with 1-2 substituents selected from the group of halo, C₁-C₄ alkyl, C₁-C₄ alkoxy, -NO₂, -NH₂, -OH and benzyloxy,
- 15 (f) benzyl, optionally substituted on the phenyl ring with one or two substituents selected from the group consisting of halo, C₁-C₄ alkyl, C₁-C₄ alkoxy and -NO₂;


20 R⁷, R⁸, R⁹, R¹⁰ are independently

- (a) H,
- (b) C₁-C₄ alkyl,
- (c) C₁-C₄ perfluoroalkyl,
- (d) C₃-C₆ cycloalkyl,
- 25 (e) phenyl, optionally substituted with one or two substituents selected from the group of halo, C₁-C₃ alkyl, C₁-C₃ alkoxy, NO₂, CF₃, NH₂, and OH,
- (f) R⁷ and R⁸ taken together are -(CH₂)_t-, or -(CH₂)_mX(CH₂)_q-,
- 30 (g) R⁹ and R¹⁰ taken together can be S, O;

13

R¹³ is

- (a) -CONHSO₂R²⁰,
(b) -NHCONHSO₂R²⁰,
(c) -NHSO₂NHCOR¹⁹,
5 (d) -PO₃H₂,
(e) -SO₃H,
(f) -SO₂NHR¹⁹,
(g) -SO₂NHCOR¹⁹,
(h) -SO₂NHCONHR¹⁹,
10 (i)



or a pharmaceutically acceptable salt thereof.

Still more preferred are compounds of the above
15 preferred scope formula (I) wherein

R⁶ is

- (a) C₁-C₇ alkyl,
(b) C₃-C₄ alkenyl,
(c) C₃-C₄ alkynyl,
20 (d) phenyl, optionally substituted with 1-2
substituents selected from the group of halo, C₁-C₄
alkyl, C₁-C₄ alkoxy, -NO₂, -NH₂, -OH and benzyloxy;


R¹³ is

- 25 (a) -CONHSO₂R²⁰,
(b) -NHCONHSO₂R²⁰,
(c) -NHSO₂NHCOR¹⁹,
(d) -SO₂NHR¹⁹,
(e) -SO₂NHCOR¹⁹,
30 (f) -SO₂NHCONHR¹⁹;

or a pharmaceutically acceptable salt thereof.

Most preferred due to their activity as angiotensin II antagonists are compounds of the more preferred scope wherein

R¹ is



or a pharmaceutically acceptable salt thereof.

Illustrative of the most preferred compounds of the invention are the following:

- 10 • N-((4'-(((4-oxo-2-propyl-8-thia-1,3-diazaspiro-((4.5))dec-1-en-3-yl-methyl))-((1,1'-biphenyl))-2-ylsulfonyl))-benzamide
- 15 • N-((4'-(((4-oxo-2-propyl-1,3-diazaspiro((4.4))non-1-en-3-yl)methyl))((1,1'-biphenyl-2-ylsulfonyl))-benzamide
- 20 • N-((4'-(((4-oxo-2-butyl-1,3-diazaspiro((4.4))non-1-en-3-yl)methyl))((1,1'-biphenyl-2-ylsulfonyl))-benzamide
- 25 • N-((4'-((4,5-dihydro-4,4-dimethyl-5-oxo-2-propyl-1H-imidazol-1-ylmethyl))((1,1'-biphenyl))-2-ylsulfonyl))-benzamide
- 30 • N-((4'-((4,5-dihydro-4,4-dimethyl-5-oxo-2-propyl-1H-imidazol-1-ylmethyl))((1,1'-biphenyl))-2-ylsulfonyl))-4-chlorobenzamide

- N-((4'-((4-oxo-2-propyl-8-thia-1,3-diazaspiro-
((4.5))dec-1-en-3-yl-methyl))-((1,1'-biphenyl))-2-
ylsulfonyl))-hexanamide
- 5 • N-((4'-((4-oxo-2-propyl-1,3-diazaspiro((2.4))hept-
1-en-3-yl)methyl))-((1,1'-biphenyl-2-ylsulfonyl))-
hexanamide
- 10 • N-((4'-((4-oxo-2-propyl-1,3-diazaspiro((4.4))non-
1-en-3-yl)methyl))-((1,1'-biphenyl-2-ylsulfonyl))-
hexanamide
- 15 • N-((4'-((4-oxo-2-propyl-1,3-diazaspiro((4.5))dec-1-
en-3-yl)methyl))-((1,1'-biphenyl-2-ylsulfonyl))-
hexanamide
- N-((4'-((4-oxo-2-butyl-1,3-diazaspiro((4.4))non-1-
en-3-yl)methyl))-((1,1'-biphenyl-2-ylsulfonyl))-
hexanamide
- 20 • N-((4'-(4,5-dihydro-4,4-dimethyl-5-oxo-2-propyl-1H-
imidazol-1-ylmethyl))-((1,1'-biphenyl))-2-
ylsulfonyl))-hexanamide
- 25 • N-((4'-(4,5-dihydro-4,4-dicyclopropyl-5-oxo-2-
propyl-1H-imidazol-1-ylmethyl))-((1,1'-biphenyl))-2-
ylsulfonyl))-hexanamide
- 30 • N-((4'-(4,5-dihydro-4,4-bis(trifluoromethyl)-5-oxo-2-
propyl-1H-imidazol-1-ylmethyl))-((1,1'-biphenyl))-2-
ylsulfonyl))-hexanamide

- N-((4'-(4,5-dihydro-4,4-dimethyl-5-oxo-2-propyl-1H-imidazol-1-ylmethyl)-3'-methyl((1,1'-biphenyl))-2-ylsulfonyl))-hexanamide
- 5 • N-((4'-(4,5-dihydro-4,4-dimethyl-5-oxo-2-propyl-1H-imidazol-1-ylmethyl)-4-propyl((1,1'-biphenyl))-2-ylsulfonyl))-hexanamide
- 10 • N-((4'-(4,5-dihydro-4,4-dimethyl-5-oxo-2-propyl-1H-imidazol-1-ylmethyl)((1,1'-biphenyl))-2-ylsulfonyl))-trifluoroacetamide
- 15 • 3,5-dihydro-5,5-dimethyl-2-propyl-3-[(2'-(N-((phenylsulfonyl)carboxamido)biphen-4-yl)methyl)]-4H-imidazol-4-one

Pharmaceutically suitable salts include both the metallic (inorganic) salts and organic salts; a list of which is given in Remington's Pharmaceutical Sciences, 20 17th Edition, page 1418 (1985). It is well known to one skilled in the art that an appropriate salt form is chosen based on physical and chemical stability, flowability, hygroscopicity, and solubility. Preferred salts of this invention for reasons cited above include 25 potassium, sodium, calcium, and ammonium salts.


Detailed Description

Synthesis

30 The compounds of formula (I) may be prepared using the reactions and techniques described in this section. The reactions are performed in solvent suitable to the reagents and materials employed and suitable for the transformation being effected. It is understood by those skilled in the art of organic synthesis that the

functionality present on the imidazole and other portions of the molecule must be consistent with the chemical transformations proposed. This will frequently necessitate judgment as to the order of synthetic steps, protecting groups required, deprotection conditions and activation of a benzylic position to enable attachment to nitrogen on the imidazole nucleus. Throughout the following section, not all compounds of formula (I) falling into a given class may necessarily be prepared by all the methods described for that class. Substituents on the starting materials may be incompatible with some of the reaction conditions required in some of the methods described. Such restrictions to the substituents which are compatible with the reaction conditions will be readily apparent to one skilled in the art and alternative methods described must then be used. The compounds of this application that have a chiral center may be resolved into the pure or partially pure optical isomers by any of the appropriate procedures known to those skilled in the art.

The compounds of formula (I) can be prepared by alkylating the alkali-metal salt of the imidazoline **1a** using appropriately protected benzyl halide, mesylate (OMs), or tosylate (OTs) derivatives **2** as shown in Scheme 1

SCHEME 1


5 The alkylation produces a mixture of the two
 regioisomers using either sodium hydride or potassium
 carbonate as base. The N¹ regioisomer is the major and
 the N³ is the minor products. These two isomers can be
 separated and purified using conventional separation
 10 techniques such as chromatography or crystallization.
 In those cases where separation of regioisomers is
 difficult by conventional techniques, the mixture can be
 transformed into suitable derivatives that can be
 separated by usual separation methods. They possess
 15 distinct physical and biological properties.

The benzyl halides of formula 2 can be prepared as
 described in European Patent Applications 400 974;
 401 030; 400,835 and references therein.

20 The starting imidazolinones are readily available
 by any number of standard methods. For example
 imidazolinone of formula 1 can be prepared as shown in
 Scheme 2. The amino nitrile 3 is readily obtainable
 from aldehydes and ketones via the Strecker Synthesis
 and various modifications thereof (R⁷ = R⁸ = CF₃, Y. V.
 25 Zeifman, N. P. Gambaryan, I. L. Knunyants, Dokl. Acad.

Nauk.S.S.S.R., 153, 1334, 1963). Treatment of the amino nitrile with triethyl amine and one equivalent of the appropriate acyl or aroyl chloride **4** in methylene chloride at room temperature overnight, gives the
5 corresponding amidonitrile **5**. Alternatively, the nitrile can be made following the procedure described in German patent disclosure DE3704100A1. The nitrile can be hydrolyzed to the diamide **6** using standard procedures such as treatment with hydrochloric acid followed by
10 ammonium hydroxide. Treatment of the diamide with 1 N sodium hydroxide as described in E. Mohr, J. Pract. Chem., 81, 49, 1910, gives the imidazolinone **1**.


Alternatively, imidazolinones of formula **1** can also be prepared as shown in Scheme 3. Treatment of the
15 amino acid **7** with tert-butyl pyrocarbonate **4** with two or more equivalents of base gives the BOC (tert-butyloxycarbonyl) protected amino acid **5**, M. Bodanszky and A. Bodanszky, The Practice of Peptide Chemistry, 1984. The protected amino amide **6** can be synthesized
20 from the active ester followed by treatment with ammonia. Deprotection using HCl gas gives the amino amide hydrochloride **7**. Treatment with two or more equivalents of base and the appropriate acyl or aroyl

SCHEME 2

5 chloride gives the diamide 6 which can be cyclized by treatment with 1 N sodium hydroxide as described above.

Likewise, compound 6 may be obtained by reacting amino acid with the requisite acid chloride by either a Schotten-Baumann procedure, or simply stirring in a solvent such as methylene chloride in the presence of base such as sodium bicarbonate, pyridine or triethyl amine followed by coupling reaction with ammonia via a variety of amide or peptide forming reactions such as DCC coupling, azide coupling, mixed anhydride synthesis or any other coupling procedure familiar to one skilled in the art.


The use of 1-amino-1-cycloalkylcarboxylic acids in the above procedure provides the imidazolinone starting materials for the preparation of the spiro-substituted imidazolinones of formula (I).

SCHEME 3

5 Imidazolinones of formula 1 can also be prepared following the procedure described in Japanese Patent disclosure JP 58055467, and the procedure by H. Lehr, J. Am. Chem. Soc., 75, 3640, 1953 and references therein.


10 Imidazolinones of formula 1 wherein R⁷ and R⁸ are both phenyl can be prepared as shown in Scheme 4 by

reaction of benzil **12** with alkyl or aryl amidine hydrochloride **13**, A. W. Cox, Org. Syn., **1**, 5, R. T. Boere, R. T. Oakley, R. W. Reed, J. Organomet. Chem., **331**, 161, 1987, in the presence of base such as 1 N sodium hydroxide, G. Rio and A. Rajon, Bull. Soc. Chim. France, 543, 1958 and references therein.

SCHEME 4

10

The imidazoline thiones of formula **15** can be prepared by treatment of the requisite alkylated imidazolinone **14** with Lawesson's reagent or phosphorus pentasulfide as described in M. P. Cava and M. I. Levinson, Tetrahedron, **41**, 5061, 1985 (Scheme 5).

SCHEME 5

20

The compounds of this invention and their preparation can be understood further by the following

examples which do not constitute a limitation of the invention. In these examples, unless otherwise indicated, all temperatures are in degrees centigrade and parts and percentages are by weight.

5

EXAMPLE 1

10 N-((4'-((4-oxo-2-propyl-8-thia-1,3-diazaspiro-((4.5))dec-1-en-3-yl-methyl))-((1,1'-biphenyl))-2-ylsulfonyl))-benzamide

PART A: Preparation of 4-Amino-4-cyanotetrahydrothio-
pyrane

15

Sodium cyanide (2.11 g, 43 mmol) was dissolved in water (40 ml). Ammonium chloride (2.53 g, 47.3 mmol) was added followed by a solution of tetrahydrothiopyran-4-one (5.0 g, 43 mmol) in methanol (40 ml). The mixture was stirred at room temperature under N₂ overnight. The mixture was diluted with H₂O and extracted with methylene chloride. The combined organic layer was washed with brine, dried over MgSO₄ and concentrated. The residue was chromatographed on silica gel eluting with hexane-ethyl acetate (1:1) to give 5.28 g of white solid (86%). MS m/e 143.0 (M+H)⁺; ¹HNMR (CDCl₃/TMS) δ 1.60-2.00(m, 4H, CH₂), 2.25(d, 2H, CH₂), 2.62-3.00(m, 4H, CH₂ and NH₂); IR(KBr, cm⁻¹) 2218.6(s, CN), 3371.3 & 3302.5(d, NH₂).

25
30

PART B: Preparation of 4-N-butyramido-4-cyanotetrahydrothiopyrane

Butyryl chloride (5.8 ml, 40.8 mmol) was added
5 dropwise to a cooled mixture of 4-Amino-4-cyano-
tetrahydrothiopyrane (5.28 g, 37.1 mmol) and triethyl
amine (5 ml) in methylene chloride (150 ml). The
mixture was stirred for 3 hours at room temperature
after which it was poured into 1N HCl (50 ml). The
10 organic layer was washed with 1N HCl (2x50 ml), 1N NaOH
(2x50 ml), dried (MgSO₄) and concentrated. The residue
was triturated with hexane to give a white solid (7.50
g, 95%). MS m/e 213, (M+H)⁺, ¹HNMR (CDCl₃/TMS) δ 0.98(t,
3H, CH₃), 1.68(m, 2H, CH₂), 1.96(m, 2H, CH₂), 2.20(t, 2H,
15 CH₂), 2.70(m, 4H, CH₂), 3.01(m, 2H, CH₂), 5.50(s, 1H,
NH).

PART C: Preparation of 4-Aminocarbonyl-4-N-butyramido-tetrahydrothiopyrane

20

4-N-butyramido-4-cyanotetrahydro-thiopyrane (7.5 g,
35.3 mmol) was dissolved in concentrated hydrochloric
acid (50 ml) at 0°C. Cold water (175 ml) was added
immediately followed by treatment with concentrated
25 ammonium hydroxide to pH 5-6. The mixture was extracted
successively with methylene chloride. The organic layer
was combined, washed with brine, dried over MgSO₄ and
concentrated to give white solid (7.0 g, 86%). MS m/e
231 (M+H)⁺, ¹HNMR (DMSO-d₆/TMS) δ 0.86(t, 3H, CH₃),
30 1.50(s, 2H, CH₂), 1.90(t, 2H, CH₂), 2.18(t, 2H, CH₂),
2.20-2.47(m, 4H, CH₂), 2.76(t, 2H, CH₂), 6.90(t, 2H,
NH₂), 7.68(s, 1H, NH).

PART D: Preparation of 4-Oxo-2-propyl-8-thia-1,3-
diazaspiro-(4,5)dec-1-ene

4-Aminocarbonyl-4-N-butylamido-tetrahydrothiopyrane
5 (7.0 g, 30.4 mmol) was heated with 1N sodium hydroxide
(50 ml) for 30 minutes. The mixture was cooled to room
temperature and some solid material was filtered off.
The filtrate was neutralized with aqueous HCl and the
white precipitate formed was filtered and dried (2.63
10 g). The aqueous layer was extracted with methylene
chloride. The combined organic layer was washed with
brine, dried and concentrated to a white solid (1.35 g).
A total of 3.98 g of product was isolated (62%). MS m/e
213 (M+H)⁺, ¹HNMR (CDCl₃/TMS) δ 1.00(t, 3H, CH₃), 1.60-
15 1.80(m, 4H, CH₂), 2.05(m, 2H, CH₂), 2.44(t, 2H, CH₂),
2.75(m, 2H, CH₂), 3.02(t, 2H, CH₂), 8.30(s, 1H, NH).

PART E: Preparation of N-[(4'-[[(4-oxo-2-propyl-8-thia-
1,3-diazaspiro-(4,5)dec-1-en-3-yl-methyl)]-N-tert-
20 butyl[(1,1'-biphenyl)]-2-sulfonamid

4-Oxo-2-propyl-8-thia-1,3-diazaspiro-(4,5)dec-1-ene
(0.83 g, 3.9 mmol) was dissolved in dimethyl formamide
(20 ml). Sodium hydride (0.18 g of 80% dispersion in
25 mineral oil) was added portionwise. The mixture was
allowed to stir at room temperature for 15 minutes. 4'-
bromomethyl-N-tert-butyl(1,1'-biphenyl)-2-sulfonamide
(1.5 g, 3.9 mmol) was added. The mixture was stirred at
room temperature under N₂ for 24 h. The solvent was
30 removed *in vacuo*. The residue was dissolved in ethyl
acetate and washed with water and brine. It was then
dried over MgSO₄, concentrated and chromatographed on
silica gel eluting with hexane-ethyl acetate (1:1) to
yield 1.65 g of white solid (82%). MS m/e 514.2 (M+H)⁺,

¹HNMR(CDCl₃/TMS) δ 0.98(t, 3H, CH₃), 0.99(s, 9H, CH₃),
1.74(m, 4H, CH₂), 2.10(m, 2H, CH₂), 2.34(t, 2H, CH₂),
2.79(m, 2H, CH₂), 3.09(t, 2H, CH₂), 3.45(s, 1H, NH),
4.73(s, 2H, CH₂Ar), 7.25(m, 3H, ArH), 7.50(m, 4H, ArH),
8.18(d, 1H, ArH).

PART F: Preparation of N-((4'-((4-oxo-2-propyl-8-thia-1,3-diazaspiro-((4.5))dec-1-en-3-yl-methyl))-(1,1'-biphenyl))-2-sulfonamide

10

N-((4'-((4-oxo-2-propyl-8-thia-1,3-diazaspiro-((4.5))dec-1-en-3-yl-methyl))-N-tert-butyl-(1,1'-biphenyl))-2-sulfonamide (1.56 g, 3.0 mmol) was refluxed with trifluoroacetic acid (10 ml) under N₂ for 2 h. The
solvent was removed *in vacuo* and the residue was dissolved in methylene chloride. The organic solution was washed with aqueous NaHCO₃ and brine. It was filtered through phase transfer paper and concentrated to an off-white solid (1.31 g, 95%). MS m/e 458.0
(M+H)⁺, ¹HNMR(CDCl₃/TMS) δ 0.97(t, 3H, CH₃), 1.60-1.82(m, 4H, CH₂), 2.10(m, 2H, CH₂), 2.36(t, 2H, CH₂), 2.78(m, 2H, CH₂), 3.09(t, 2H, CH₂), 4.21(s, 2H, NH₂), 4.73(s, 2H, CH₂Ar), 7.20(d, 2H, ArH), 7.30(d, 1H, ArH), 7.42-7.63(m, 4H, ArH), 8.17(d, 1H, ArH).

25

PART G: Preparation of N-((4'-((4-oxo-2-propyl-8-thia-1,3-diazaspiro-((4.5))dec-1-en-3-yl-methyl))-((1,1'-biphenyl))-2-ylsulfonyl))-benzamide

30

1,1'-Carbonyl diimidazole (1.26 g, 7.8 mmol) and benzoic acid (0.96 g, 7.8 mmol) was refluxed with tetrahydrofuran (30 ml) under N₂ for 2 h. The mixture was cooled to room temperature, and a solution of N-((4'-((4-oxo-2-propyl-8-thia-1,3-diazaspiro-

((4.5))dec-1-en-3-yl-methyl))((1,1'-biphenyl))-2-sulfonamide (1.20 g, 2.6 mmol) and 1,8-diazabicyclo[5.4.0]undec-7-ene (1.2 ml, 7.8 mmol) in THF (30 ml) was added. The reaction mixture was then
5 refluxed for 1.5 h. The mixture was cooled and poured into 20 ml of 25% aqueous citric acid. It was extracted with ethyl acetate. The combined organic solution was washed with brine, dried over MgSO₄ and concentrated. the residue was chromatographed on silica gel eluting
10 with 5% methanol in methylene chloride to yield 1.3 g of the desired product (91%). M.P. 105-108°C, MS m/e 561 (M+H)⁺, ¹HNMR(CDCl₃/TMS) δ 0.98(t, 3H, CH₃), 1.60-1.80(m, 4H, CH₂), 2.08(t, 2H, CH₂), 2.34(t, 2H, CH₂), 2.78(m, 2H, CH₂), 3.08(t, 2H, CH₂), 4.65(s, 2H, CH₂Ar), 6.99(d, 2H, ArH),
15 7.23(d, 1H, ArH), 7.42(m, 4H, ArH), 7.60(m, 4H, ArH), 8.40(d, 1H, ArH).

EXAMPLE 2

N-((4'-((4-oxo-2-propyl-1,3-diazaspiro((4.4))non-1-en-3-yl)methyl))((1,1'-biphenyl-2-yl)sulfonyl))-benzamide
20

PART A: Preparation of 4-Oxo-2-propyl-1,3-diazaspiro-(4,4)non-1-ene

25 1-Amino-1-cyclopentane carboxylic acid methyl ester (10.1 g, 70.6 mmol), ethyl butanimidate hydrochloride (12.7 g, 84.7 mmol) and triethyl amine (17 ml) was refluxed in benzene (50 ml) under N₂ overnight. The solvent was removed *in vacuo* and the residue was
30 dissolved in methylene chloride, washed with water and brine, and concentrated. The crude product mixture was chromatographed on silica gel eluting with ethyl acetate to give 8.47 g of colorless oil (67%). MS m/e 181.1


(M+H)⁺, ¹HNMR(CDCl₃/TMS) δ 1.00(t, 3H, CH₃), 1.50-2.20(m, 10H, CH₂), 2.41(t, 2H, CH₂), 9.09(br.s, 1H, NH).

PART B: N-((4'-((4-oxo-2-propyl-1,3-diazaspiro-
5 ((4,4))non-1-en-3-yl)methyl))((1,1'-biphenyl-2-
ylsulfonyl))-bebzamide

The titled compound was prepared from 4-Oxo-2-propyl-1,3-diazaspiro-(4,4)non-1-ene and 4'-bromomethyl-
10 N-tert-butyl(1,1'-biphenyl)-2-sulfonamide according to the procedures described in Part E, F and G of Example 1. M.P. 118-125°C, MS m/e 529 (M+H)⁺,
¹HNMR(DMSO-d₆/TMS) δ 0.89(t, 3H, CH₃), 1.50-1.92(m, 10H, CH₂), 2.34(t, 2H, CH₂), 4.68(s, 2H, CH₂Ar), 7.00(d, 2H,
15 ArH), 7.10-7.62(m, 10H, ArH), 8.10(d, 1H, ArH).

Compounds 1-630 in Table 1 can be prepared by the procedures described in Examples 1 and 2 employing the appropriately substituted imidazolines and benzyl halides or mesylates.

TABLE 1



5	EX.	R ⁶	R ⁷	R ⁸	R ⁹ R ¹⁰	R ¹³	R ² , R ³	MS (M+H) ⁺
	1	n-Pr	-(CH ₂) ₂ S(CH) ₂ -		O	-SO ₂ NHCOC ₆ H ₅	H, H	561
	2	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCOC ₆ H ₅	H, H	529
	3	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (n-C ₄ H ₉)	H, H	
10	4	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCOC ₆ H ₅	H, H	543
	5	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCOC ₆ H ₅	H, H	504
	6	n-Pr	CH ₃	CH ₃	O	-CONHSO ₂ CH ₂ C ₆ H ₅	H, H	
	7	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCO (n-C ₅ H ₁₁) x	H, H	
	8	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, H	
15	9	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H	
	10	n-Pr	CH ₃	CH ₃	O	-CH ₂ CO ₂ H	H, H	
	11	n-Pr	CH ₃	CH ₃	O	-C(CF ₃) ₂ OH	H, H	
	12	n-Pr	CH ₃	CH ₃	O	-CONHNHSO ₂ CF ₃	H, H	
	13	n-Pr	CH ₃	CH ₃	O	-CONHOCH ₃	H, H	
20	14	n-Pr	CH ₃	CH ₃	O	-CONHSO ₂ C ₆ H ₅	H, H	
	15	n-Pr	CH ₃	CH ₃	O	-PO ₃ H ₂	H, H	
	16	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCOC ₆ H ₅	CH ₃ , H	
	17	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	CH ₃ , H	
	18	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCO (cy-C ₃ H ₅)	CH ₃ , H	
25	19	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	CH ₃ , H	
	20	n-Pr	CH ₃	CH ₃	O	-SO ₃ H	CH ₃ , H	
	21	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCONH (n-Bu)	CH ₃ , H	
	22	n-Pr	CH ₃	CH ₃	O	-SO ₂ NH (C ₅ NH ₄)	CH ₃ , H	

30

	EX.	R ⁶	R ⁷	R ⁸	R ⁹ R ¹⁰	R ¹³	R ² , R ³
	23	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCONH (n-C ₅ H ₁₁)	CH ₃ , H
	24	n-Pr	CH ₃	CH ₃	O	-NH ₂ SO ₂ NHCO (i-C ₅ H ₁₁)	CH ₃ , H
	25	n-Pr	CH ₃	CH ₃	O	-NH ₂ SO ₂ NHCO (cy-C ₃ H ₅)	CH ₃ , H
5	26	n-Pr	CH ₃	CH ₃	O	-NH ₂ SO ₂ NHCOCH ₂ C ₆ H ₅	CH ₃ , H
	27	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCOC ₆ H ₅	Cl, H
	28	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	Cl, H
	29	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCO (cy-C ₃ H ₅)	Cl, H
	30	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	Cl, H
10	31	n-Pr	CH ₃	CH ₃	O	-OPO ₃ H ₂	Cl, H
	32	n-Pr	CH ₃	CH ₃	O	-PO (OH) CH ₂ C ₆ H ₅	Cl, H
	33	n-Pr	CH ₃	CH ₃	O	-OSO ₃ H	Cl, H
	34	n-Pr	CH ₃	CH ₃	O	-NHPO ₃ H ₂	Cl, H
	35	n-Pr	CH ₃	CH ₃	O	-SO ₂ NH ₂	Cl, H
15	36	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHC ₂ H ₅	Cl, H
	37	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHC ₁₀ H ₇	Cl, H
	38	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCOC ₆ H ₅	F, H
	39	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	F, H
	40	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCO (cy-C ₃ H ₅)	F, H
20	41	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	F, H
	42	n-Pr	CH ₃	CH ₃	O	-NH ₂ SO ₂ NHCO (n-Bu)	F, H
	43	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCONH (n-Bu)	F, H
	44	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCO (i-Bu)	F, H
	45	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCO (4-HOC ₆ H ₅)	F, H
25	46	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	F, H
	47	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCO (4-C ₅ NH ₄)	F, H
	48	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCONHCH ₂ C ₆ H ₅	F, H
	49	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCOC ₆ H ₅	H, n-Pr
	50	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, n-Pr
30	51	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, n-Pr
	52	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, n-Pr
	53	n-Pr	CH ₃	CH ₃	O	-CH ₂ CO ₂ H	H, n-Pr
	54	n-Pr	CH ₃	CH ₃	O	-C (CF ₃) ₂ OH	H, n-Pr
	55	n-Pr	CH ₃	CH ₃	O	-CONHNH ₂ SO ₂ CF ₃	H, n-Pr

	EX.	R ⁶	R ⁷	R ⁸	R ⁹ R ¹⁰	R ¹³	R ² , R ³
	56	n-Pr	CH ₃	CH ₃	O	-CONHOH	H, n-Pr
	57	n-Pr	CH ₃	CH ₃	O	-CONHOCH ₃	H, n-Pr
	58	n-Pr	CH ₃	CH ₃	O	-CONHOCH ₂ C ₆ H ₅	H, n-Pr
5	59	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCOC ₆ H ₅	Cl, n-Pr
	60	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	Cl, n-Pr
	61	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCO (cy-C ₃ H ₅)	F, n-Pr
	62	n-Pr	CH ₃	CH ₃	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	F, n-Pr
	63	n-Pr	CH ₃	CH ₃	O	-CONHSO ₂ (4-ClC ₆ H ₄)	Cl, n-Pr
10	64	n-Pr	CH ₃	CH ₃	O	-CONHSO ₂ C ₂ F ₅	F, n-Pr
	65	n-Pr	CH ₃	CH ₃	O	-CONHSO ₂ C ₂ H ₄ OH	Cl, n-Pr
	66	n-Pr	CH ₃	CH ₃	O	-CONHSO ₂ C ₂ H ₄ CO ₂ H	F, n-Pr
	67	n-Pr	CH ₃	CH ₃	O	-CONHSO ₂ C ₂ H ₄ NH ₂	F, n-Pr
	68	n-Pr	CH ₃	CH ₃	O	-CONHSO ₂ NH ₂	Cl, n-Pr
15	69	n-Bu	CH ₃	CH ₃	O	-SO ₂ NHCOC ₆ H ₅	H, H
	70	n-Bu	CH ₃	CH ₃	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	71	n-Bu	CH ₃	CH ₃	O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, H
	72	n-Bu	CH ₃	CH ₃	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	73	n-Bu	CH ₃	CH ₃	O	-CONHSO ₂ NHC ₂ H ₅	H, H
20	74	n-Bu	CH ₃	CH ₃	O	-CONHSO ₂ NHC ₆ H ₅	H, H
	75	n-Bu	CH ₃	CH ₃	O	-CONHSO ₂ NHCH ₂ C ₆ H ₅	H, H
	76	n-Bu	CH ₃	CH ₃	O	-CONHSO ₂ NH (4-C ₅ NH ₄)	H, H
	77	n-Bu	CH ₃	CH ₃	O	-C (OH) CH ₃ PO ₃ H ₂	H, H
	78	n-Bu	CH ₃	CH ₃	O	-C (OH) HPO ₃ H ₂	H, H
25	79	n-Bu	CH ₃	CH ₃	O	-SO ₂ NHCOC ₆ H ₅	CH ₃ , H
	80	n-Bu	CH ₃	CH ₃	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	CH ₃ , H
	81	n-Bu	CH ₃	CH ₃	O	-SO ₂ NHCO (cy-C ₃ H ₅)	CH ₃ , H
	82	n-Bu	CH ₃	CH ₃	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	CH ₃ , H
	83	n-Bu	CH ₃	CH ₃	O	-NHCONHSO ₂ C ₂ H ₅	CH ₃ , H
30	84	n-Bu	CH ₃	CH ₃	O	-NHCONHSO ₂ (i-Bu)	CH ₃ , H
	85	n-Bu	CH ₃	CH ₃	O	-NHCONHSO ₂ (n-C ₅ H ₁₁)	CH ₃ , H
	86	n-Bu	CH ₃	CH ₃	O	-SO ₂ NHCONH (n-C ₅ H ₁₁)	CH ₃ , H
	87	n-Bu	CH ₃	CH ₃	O	-SO ₂ NHCO (i-C ₅ H ₁₁)	CH ₃ , H
	88	n-Bu	CH ₃	CH ₃	O	-SO ₂ NHCO (cy-C ₃ H ₅)	CH ₃ , H

	EX.	R ⁶	R ⁷	R ⁸	R ⁹ R ¹⁰	R ¹³	R ² , R ³
	89	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHCOCH ₂ C ₆ H ₅	CH ₃ , H
	90	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHCOC ₆ H ₅	Cl, H
	91	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHCO (n-C ₅ H ₁₁)	Cl, H
5	92	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHCO (cy-C ₃ H ₅)	Cl, H
	93	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHCOCH ₂ C ₆ H ₅	Cl, H
	94	n-Bu	CH ₃	CH ₃	0	-NHPO ₃ H ₂	Cl, H
	95	n-Bu	CH ₃	CH ₃	0	-NH ₂ SO ₂ NHCO (Cy-C ₃ H ₅)	Cl, H
	96	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHCONH (i-Bu)	Cl, H
10	97	n-Bu	CH ₃	CH ₃	0	-PO (OH) (n-C ₅ H ₁₁)	Cl, H
	98	n-Bu	CH ₃	CH ₃	0	-PO (OH) (i-C ₅ H ₁₁)	Cl, H
	99	n-Bu	CH ₃	CH ₃	0	-PO (OH) C ₃ H ₇	Cl, H
	100	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHCH ₂ C ₆ H ₅	Cl, H
	101	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHC ₆ H ₅	F, H
15	102	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHCO (n-C ₅ H ₁₁)	F, H
	103	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHCO (cy-C ₃ H ₅)	F, H
	104	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHCOCH ₂ C ₆ H ₅	F, H
	105	n-Bu	CH ₃	CH ₃	0	-SO ₂ NH (n-Bu)	F, H
	106	n-Bu	CH ₃	CH ₃	0	-SO ₂ NH (i-Bu)	F, H
20	107	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHCONH (i-Bu)	F, H
	108	n-Bu	CH ₃	CH ₃	0	-SO ₂ NH (n-C ₅ H ₁₁)	F, H
	109	n-Bu	CH ₃	CH ₃	0	-SO ₂ NH (i-C ₅ H ₁₁)	F, H
	110	n-Bu	CH ₃	CH ₃	0	-NH ₂ SO ₂ NHCO (cy-C ₃ H ₅)	F, H
	111	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHCOCH ₂ C ₆ H ₅	F, H
25	112	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHCOC ₆ H ₅	H, n-Pr
	113	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, n-Pr
	114	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHCO (cy-C ₃ H ₅)	H, n-Pr
	115	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, n-Pr
	116	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHCO (n-Bu)	H, n-Pr
30	117	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHCONH (n-Bu)	H, n-Pr
	118	n-Bu	CH ₃	CH ₃	0	-SO ₂ NHCONH (i-Bu)	H, n-Pr
	119	n-Bu	CH ₃	CH ₃	0	-SO ₂ NH (n-C ₅ H ₁₁)	H, n-Pr
	120	n-Bu	CH ₃	CH ₃	0	-SO ₂ NH (i-C ₅ H ₁₁)	H, n-Pr
	121	n-Bu	CH ₃	CH ₃	0	-SO ₂ NH (cy-C ₃ H ₅)	H, n-Pr

	EX.	R ⁶	R ⁷	R ⁸	R ⁹ R ¹⁰	R ¹³	R ² , R ³
	122	n-Bu	CH ₃	CH ₃	O	-SO ₂ NHCH ₂ C ₆ H ₅	H, n-Pr
	123	n-Bu	CH ₃	CH ₃	O	-SO ₂ NHCOC ₆ H ₅	Cl, n-Pr
	124	n-Bu	CH ₃	CH ₃	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	Cl, n-Pr
5	125	n-Bu	CH ₃	CH ₃	O	-SO ₂ NHCO (cy-C ₃ H ₅)	F, n-Pr
	126	n-Bu	CH ₃	CH ₃	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	F, n-Pr
	127	n-Bu	CH ₃	CH ₃	O	-SO ₂ NH (n-Bu)	Cl, n-Pr
	128	n-Bu	CH ₃	CH ₃	O	-SO ₂ NH (i-Bu)	F, n-Pr
	129	n-Bu	CH ₃	CH ₃	O	-CONHSO ₂ (n-C ₅ H ₁₁)	Cl, n-Pr
10	130	n-Bu	CH ₃	CH ₃	O	-CONHSO ₂ (i-C ₅ H ₁₁)	F, n-Pr
	131	n-Bu	CH ₃	CH ₃	O	-CONHSO ₂ (cy-C ₃ H ₅)	F, n-Pr
	132	n-Bu	CH ₃	CH ₃	O	-CONHSO ₂ CH ₂ C ₆ H ₅	Cl, n-Pr
	133	Ph	CH ₃	CH ₃	O	-SO ₂ NHCOC ₆ H ₅	H, H
	134	Ph	CH ₃	CH ₃	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, H
15	135	Ph	CH ₃	CH ₃	O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, H
	136	p-F-Ph	CH ₃	CH ₃	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	137	p-F-Ph	CH ₃	CH ₃	O	-SO ₂ NHCO (n-Bu)	H, H
	138	Ph	CH ₃	CH ₃	O	-SO ₂ NHCO (i-Bu)	H, H
	139	iPr	CH ₃	CH ₃	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, H
20	140	Ph	CH ₃	CH ₃	O	-SO ₂ NHCO (i-C ₅ H ₁₁)	H, H
	141	Ph	CH ₃	CH ₃	O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, H
	142	Ph	CH ₃	CH ₃	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	143	n-Pr	-(CH ₂) ₄ -		O	-CONHSO ₂ CH ₂ C ₆ H ₅	H, H
	144	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, H
25	145	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, H
	146	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	147	n-Pr	-(CH ₂) ₄ -		O	-CH ₂ CO ₂ H	H, H
	148	n-Pr	-(CH ₂) ₄ -		O	-C(CF ₃) ₂ OH	H, H
	149	n-Pr	-(CH ₂) ₄ -		O	-CONHNHSO ₂ CF ₃	H, H
30	150	n-Pr	-(CH ₂) ₄ -		O	-CONHOCH ₂ C ₆ H ₅	H, H
	151	n-Pr	-(CH ₂) ₄ -		O	-CONHOCH ₃	H, H
	152	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCOC ₆ H ₅	CH ₃ , H
	153	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (n-C ₅ H ₁₁)	CH ₃ , H
	154	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (cy-C ₃ H ₅)	CH ₃ , H

	EX.	R ⁶	R ⁷	R ⁸	R ⁹ R ¹⁰	R ¹³	R ² , R ³
	155	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCOCH ₂ C ₆ H ₅	CH ₃ , H
	156	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (n-Bu)	CH ₃ , H
	157	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCONH (n-Bu)	CH ₃ , H
5	158	n-Pr	-(CH ₂) ₄ -		O	-CONHSO ₂ (i-Bu)	CH ₃ , H
	159	n-Pr	-(CH ₂) ₄ -		O	-CONHSO ₂ (n-C ₅ H ₁₁)	CH ₃ , H
	160	n-Pr	-(CH ₂) ₄ -		O	-CONHSO ₂ (i-C ₅ H ₁₁)	CH ₃ , H
	161	n-Pr	-(CH ₂) ₄ -		O	-CONHSO ₂ (cy-C ₃ H ₅)	CH ₃ , H
	162	n-Pr	-(CH ₂) ₄ -		O	-CONHSO ₂ CH ₂ C ₆ H ₅	CH ₃ , H
10	163	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCOC ₆ H ₅	Cl, H
	164	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (n-C ₅ H ₁₁)	Cl, H
	165	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (cy-C ₃ H ₅)	Cl, H
	166	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCOCH ₂ C ₆ H ₅	Cl, H
	167	n-Pr	-(CH ₂) ₄ -		O	-CONHSO ₂ NH (n-Bu)	Cl, H
15	168	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCONH (n-Bu)	Cl, H
	169	n-Pr	-(CH ₂) ₄ -		O	-NHCONHSO ₂ (i-Bu)	Cl, H
	170	n-Pr	-(CH ₂) ₄ -		O	-NHCONHSO ₂ (n-C ₅ H ₁₁)	Cl, H
	171	n-Pr	-(CH ₂) ₄ -		O	-NHCONHSO ₂ (i-C ₅ H ₁₁)	Cl, H
	172	n-Pr	-(CH ₂) ₄ -		O	-NHCONHSO ₂ (cy-C ₃ H ₅)	Cl, H
20	173	n-Pr	-(CH ₂) ₄ -		O	-NHCONHSO ₂ CH ₂ C ₆ H ₅	Cl, H
	174	n-Pr	-(CH ₂) ₄ -		O	-OP ₃ H ₂	F, H
	175	n-Pr	-(CH ₂) ₄ -		O	-OSO ₃ H	F, H
	176	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (cy-C ₃ H ₅)	F, H
	177	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCOCH ₂ C ₆ H ₅	F, H
25	178	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (n-Bu)	F, H
	179	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCONH (n-Bu)	F, H
	180	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (i-Bu)	F, H
	181	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (n-C ₅ H ₁₁)	F, H
	182	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCONH (i-C ₅ H ₁₁)	F, H
30	183	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCONH (cy-C ₃ H ₅)	F, H
	184	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCH ₂ C ₆ H ₅	F, H
	185	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCOC ₆ H ₅	H, n-Pr
	186	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, n-Pr
	187	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, n-Pr

	EX.	R ⁶	R ⁷	R ⁸	R ⁹ R ¹⁰	R ¹³	R ² , R ³
	188	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, n-Pr
	189	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (n-Bu)	H, n-Pr
	190	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCONH (n-Bu)	H, n-Pr
5	191	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (i-Bu)	H, n-Pr
	192	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NH (n-C ₅ H ₁₁)	H, n-Pr
	193	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NH (i-C ₅ H ₁₁)	H, n-Pr
	194	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NH (cy-C ₃ H ₅)	H, n-Pr
	195	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCH ₂ C ₆ H ₅	H, n-Pr
10	196	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCOC ₆ H ₅	Cl, n-Pr
	197	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (n-C ₅ H ₁₁)	Cl, n-Pr
	198	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (cy-C ₃ H ₅)	F, n-Pr
	199	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCOCH ₂ C ₆ H ₅	F, n-Pr
	200	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (n-Bu)	Cl, n-Pr
15	201	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCOC ₂ H ₅	F, n-Pr
	202	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCOCH ₃	Cl, n-Pr
	203	n-Pr	-(CH ₂) ₄ -		O	-SO ₂ NHCO (i-C ₅ H ₁₁)	F, n-Pr
	204	n-Pr	-(CH ₂) ₄ -		O	-CH ₂ CO ₂ H	F, n-Pr
	205	n-Pr	-(CH ₂) ₄ -		O	-CONHNH ₂ SO ₂ CF ₃	Cl, n-Pr
20	206	n-Bu	-(CH ₂) ₄ -		O	-CONHSO ₂ (n-C ₅ H ₁₁)	H, H
	207	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, H
	208	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	209	n-Bu	-(CH ₂) ₄ -		O	-CONHSO ₂ (n-Bu)	H, H
	210	n-Bu	-(CH ₂) ₄ -		O	-CONHSO ₂ (i-Bu)	H, H
25	211	n-Bu	-(CH ₂) ₄ -		O	-CONHSO ₂ (n-C ₅ H ₁₁)	H, H
	212	n-Bu	-(CH ₂) ₄ -		O	-CONHSO ₂ (i-C ₅ H ₁₁)	H, H
	213	n-Bu	-(CH ₂) ₄ -		O	-CONHSO ₂ (cy-C ₃ H ₅)	H, H
	214	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	215	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCOC ₆ H ₅	CH ₃ , H
30	216	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCO (n-C ₅ H ₁₁)	CH ₃ , H
	217	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCO (cy-C ₃ H ₅)	CH ₃ , H
	218	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCOCH ₂ C ₆ H ₅	CH ₃ , H
	219	n-Bu	-(CH ₂) ₄ -		O	-NHCONHSO ₂ (n-Bu)	CH ₃ , H
	220	n-Bu	-(CH ₂) ₄ -		O	-NHCONHSO ₂ (i-Bu)	CH ₃ , H

EX.	R ⁶	R ⁷	R ⁸	R ⁹ R ¹⁰	R ¹³	R ² , R ³
221	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCONH(i-Bu)	CH ₃ , H
222	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NH(n-C ₅ H ₁₁)	CH ₃ , H
223	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NH(i-C ₅ H ₁₁)	CH ₃ , H
5 224	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NH ₂ (cy-C ₃ H ₅)	CH ₃ , H
225	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCH ₂ C ₆ H ₅	CH ₃ , H
226	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCOC ₆ H ₅	Cl, H
227	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCO(n-C ₅ H ₁₁)	Cl, H
228	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCO(cy-C ₃ H ₅)	Cl, H
10 229	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCOCH ₂ C ₆ H ₅	Cl, H
230	n-Bu	-(CH ₂) ₄ -		O	-NH ₂ SO ₂ NHCO(n-Bu)	Cl, H
231	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCONH(n-Bu)	Cl, H
232	n-Bu	-(CH ₂) ₄ -		O	-NH ₂ SO ₂ NHCO(i-Bu)	Cl, H
233	n-Bu	-(CH ₂) ₄ -		O	-NH ₂ SO ₂ NHCO(n-C ₅ H ₁₁)	Cl, H
15 234	n-Bu	-(CH ₂) ₄ -		O	-NH ₂ SO ₂ NHCO(i-C ₅ H ₁₁)	Cl, H
235	n-Bu	-(CH ₂) ₄ -		O	-CH ₂ (5-Tetrazoyl)	Cl, H
236	n-Bu	-(CH ₂) ₄ -		O	-CONH(5-Tetrazoyl)	Cl, H
237	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCOC ₆ H ₅	F, H
238	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCO(n-C ₅ H ₁₁)	F, H
20 239	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCO(cy-C ₃ H ₅)	F, H
240	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCOCH ₂ C ₆ H ₅	F, H
241	n-Bu	-(CH ₂) ₄ -		O	-NH ₂ SO ₂ NHCO(n-Bu)	F, H
242	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCONH(n-Bu)	F, H
243	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCO(i-Bu)	F, H
25 244	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NH(n-C ₅ H ₁₁)	F, H
245	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NH(i-C ₅ H ₁₁)	F, H
246	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NH(cy-C ₃ H ₅)	F, H
247	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCH ₂ C ₆ H ₅	F, H
248	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCOC ₆ H ₅	H, n-Pr
30 249	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCO(n-C ₅ H ₁₁)	H, n-Pr
250	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCO(cy-C ₃ H ₅)	H, n-Pr
251	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, n-Pr
252	n-Bu	-(CH ₂) ₄ -		O	-CONHSO ₂ NHC ₂ H ₄ Cl	H, n-Pr
253	n-Bu	-(CH ₂) ₄ -		O	-CONHSO ₂ (i-Bu)	H, n-Pr

	EX.	R ⁶	R ⁷	R ⁸	R ⁹ R ¹⁰	R ¹³	R ² , R ³
	254	n-Bu	-(CH ₂) ₄ -		O	-CONHSO ₂ (n-C ₅ H ₁₁)	H, n-Pr
	255	n-Bu	-(CH ₂) ₄ -		O	-CONHSO ₂ (i-C ₅ H ₁₁)	H, n-Pr
	256	n-Bu	-(CH ₂) ₄ -		O	-CONHSO ₂ (cy-C ₃ H ₅)	H, n-Pr
5	257	n-Bu	-(CH ₂) ₄ -		O	-CONHSO ₂ CH ₂ C ₆ H ₅	H, n-Pr
	258	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCOC ₆ H ₅	Cl, n-Pr
	259	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCO (n-C ₅ H ₁₁)	Cl, n-Pr
	260	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCO (cy-C ₃ H ₅)	F, n-Pr
	261	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCOCH ₂ C ₆ H ₅	F, n-Pr
10	262	n-Bu	-(CH ₂) ₄ -		O	-NHSO ₂ NHCO (n-Bu)	Cl, n-Pr
	263	n-Bu	-(CH ₂) ₄ -		O	-NHSO ₂ NHCO (i-Bu)	F, n-Pr
	264	n-Bu	-(CH ₂) ₄ -		O	-NHSO ₂ NHCO (n-C ₅ H ₁₁)	Cl, n-Pr
	265	n-Bu	-(CH ₂) ₄ -		O	-NHSO ₂ NHCO (i-C ₅ H ₁₁)	F, n-Pr
	266	n-Bu	-(CH ₂) ₄ -		O	-SO ₂ NHCONH (i-C ₅ H ₁₁)	F, n-Pr
15	267	n-Bu	-(CH ₂) ₄ -		O	-NHSO ₂ NHCO (cy-C ₃ H ₅)	F, n-Pr
	268	n-Bu	-(CH ₂) ₄ -		O	-NHSO ₂ NHCOCH ₂ C ₆ H ₅	Cl, n-Pr
	269	n-Pr	-(CH ₂) ₅ -		O	-CONHSO ₂ CH ₂ C ₆ H ₅	H, H
	270	n-Pr	-(CH ₂) ₅ -		O	-SO ₂ NHCOC ₆ H ₅	H, H
	271	n-Pr	-(CH ₂) ₅ -		O	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, H
20	272	n-Pr	-(CH ₂) ₅ -		O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, H
	273	n-Pr	-(CH ₂) ₅ -		O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	280	n-Pr	-(CH ₂) ₅ -		O	-SO ₂ NHCOC ₆ H ₅	CH ₃ , H
	281	n-Pr	-(CH ₂) ₅ -		O	-SO ₂ NHCO (n-C ₅ H ₁₁)	CH ₃ , H
	282	n-Pr	-(CH ₂) ₅ -		O	-SO ₂ NHCO (cy-C ₃ H ₅)	CH ₃ , H
25	283	n-Pr	-(CH ₂) ₅ -		O	-SO ₂ NHCOCH ₂ C ₆ H ₅	CH ₃ , H
	284	n-Pr	-(CH ₂) ₅ -		O	-NHSO ₂ NHCO (n-Bu)	CH ₃ , H
	285	n-Pr	-(CH ₂) ₅ -		O	-SO ₂ NHCONH (n-Bu)	CH ₃ , H
	286	n-Pr	-(CH ₂) ₅ -		O	-NHSO ₂ NHCO (i-Bu)	CH ₃ , H
	287	n-Pr	-(CH ₂) ₅ -		O	-NHSO ₂ NHCO (n-C ₅ H ₁₁)	CH ₃ , H
30	288	n-Pr	-(CH ₂) ₅ -		O	-NHSO ₂ NHCO (i-C ₅ H ₁₁)	CH ₃ , H
	289	n-Pr	-(CH ₂) ₅ -		O	-NHSO ₂ NHCO (cy-C ₃ H ₅)	CH ₃ , H
	290	n-Pr	-(CH ₂) ₅ -		O	-NHSO ₂ NHCOCH ₂ C ₆ H ₅	CH ₃ , H
	291	n-Pr	-(CH ₂) ₅ -		O	-SO ₂ NHCOC ₆ H ₅	Cl, H
	292	n-Pr	-(CH ₂) ₅ -		O	-SO ₂ NHCO (n-C ₅ H ₁₁)	Cl, H

	EX.	R ⁶	R ⁷	R ⁸	R ⁹	R ¹⁰	R ¹³	R ² , R ³
	293	n-Pr	-(CH ₂) ₅ -		O		-SO ₂ NHCO (cy-C ₃ H ₅)	Cl, H
	294	n-Pr	-(CH ₂) ₅ -		O		-SO ₂ NHCOCH ₂ C ₆ H ₅	Cl, H
	295	n-Pr	-(CH ₂) ₅ -		O		-NHSO ₂ NHCO (n-Bu)	Cl, H
5	296	n-Pr	-(CH ₂) ₅ -		O		-NHSO ₂ NHCO (n-Bu)	Cl, H
	297	n-Pr	-(CH ₂) ₅ -		O		-SO ₂ NHCONH (i-Bu)	Cl, H
	298	n-Pr	-(CH ₂) ₅ -		O		-SO ₂ NHCO (n-C ₅ H ₁₁)	Cl, H
	299	n-Pr	-(CH ₂) ₅ -		O		-NHSO ₂ NHCO (i-C ₅ H ₁₁)	Cl, H
	300	n-Pr	-(CH ₂) ₅ -		O		-SO ₂ NH (cy-C ₃ H ₅)	Cl, H
10	301	n-Pr	-(CH ₂) ₅ -		O		-SO ₂ NHCH ₂ C ₆ H ₅	Cl, H
	302	n-Pr	-(CH ₂) ₅ -		O		-SO ₂ NHCOC ₆ H ₅	F, H
	303	n-Pr	-(CH ₂) ₅ -		O		-SO ₂ NHCO (n-C ₅ H ₁₁)	F, H
	304	n-Pr	-(CH ₂) ₅ -		O		-SO ₂ NHCO (cy-C ₃ H ₅)	F, H
	305	n-Pr	-(CH ₂) ₅ -		O		-SO ₂ NHCOCH ₂ C ₆ H ₅	F, H
15	306	n-Pr	-(CH ₂) ₅ -		O		-SO ₂ NHCO (n-Bu)	F, H
	307	n-Pr	-(CH ₂) ₅ -		O		-SO ₂ NHCONH (n-Bu)	F, H
	308	n-Pr	-(CH ₂) ₅ -		O		-NHSO ₂ NHCO (i-Bu)	F, H
	309	n-Pr	-(CH ₂) ₅ -		O		-NHSO ₂ NHCO (n-C ₅ H ₁₁)	F, H
	310	n-Pr	-(CH ₂) ₅ -		O		-NHSO ₂ NHCO (i-C ₅ H ₁₁)	F, H
20	311	n-Pr	-(CH ₂) ₅ -		O		-NHSO ₂ NHCO (cy-C ₃ H ₅)	F, H
	312	n-Pr	-(CH ₂) ₅ -		O		-NHSO ₂ NHCOCH ₂ C ₆ H ₅	F, H
	313	n-Pr	-(CH ₂) ₅ -		O		-SO ₂ NHCOC ₆ H ₅	H, n-Pr
	314	n-Pr	-(CH ₂) ₅ -		O		-SO ₂ NHCO (n-C ₅ H ₁₁)	H, n-Pr
	315	n-Pr	-(CH ₂) ₅ -		O		-SO ₂ NHCO (cy-C ₃ H ₅)	H, n-Pr
25	316	n-Pr	-(CH ₂) ₅ -		O		-SO ₂ NHCOCH ₂ C ₆ H ₅	H, n-Pr
	317	n-Pr	-(CH ₂) ₅ -		O		-NHSO ₂ NHCO (n-Bu)	H, n-Pr
	318	n-Pr	-(CH ₂) ₅ -		O		-SO ₂ NHCONH (n-Bu)	H, n-Pr
	319	n-Pr	-(CH ₂) ₅ -		O		-NHSO ₂ NHCO (i-Bu)	H, n-Pr
	320	n-Pr	-(CH ₂) ₅ -		O		-NHSO ₂ NHCO (n-C ₅ H ₁₁)	H, n-Pr
30	321	n-Pr	-(CH ₂) ₅ -		O		-NHSO ₂ NHCO (i-C ₅ H ₁₁)	H, n-Pr
	322	n-Pr	-(CH ₂) ₅ -		O		-NHSO ₂ NHCO (cy-C ₃ H ₅)	H, n-Pr
	323	n-Pr	-(CH ₂) ₅ -		O		-NHSO ₂ NHCOCH ₂ C ₆ H ₅	H, n-Pr
	324	n-Pr	-(CH ₂) ₅ -		O		-SO ₂ NHCOC ₆ H ₅	Cl, n-Pr
	325	n-Pr	-(CH ₂) ₅ -		O		-SO ₂ NHCO (n-C ₅ H ₁₁)	Cl, n-Pr

	EX.	R ⁶	R ⁷	R ⁸	R ⁹ R ¹⁰	R ¹³	R ² , R ³
	326	n-Pr	-	(CH ₂) ₅ -	O	-SO ₂ NHCO (cy-C ₃ H ₅)	F, n-Pr
	327	n-Pr	-	(CH ₂) ₅ -	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	F, n-Pr
	328	n-Pr	-	(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (n-Bu)	Cl, n-Pr
5	329	n-Pr	-	(CH ₂) ₅ -	O	-SO ₂ NHCONH (n-Bu)	Cl, n-Pr
	330	n-Pr	-	(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (i-Bu)	F, n-Pr
	331	n-Pr	-	(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (n-C ₅ H ₁₁)	Cl, n-Pr
	332	n-Pr	-	(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (i-C ₅ H ₁₁)	F, n-Pr
	333	n-Pr	-	(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (cy-C ₃ H ₅)	F, n-Pr
10	334	n-Pr	-	(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCOCH ₂ C ₆ H ₅	Cl, n-Pr
	335	n-Bu	-	(CH ₂) ₅ -	O	-SO ₂ NHCOC ₆ H ₅	H, H
	336	n-Bu	-	(CH ₂) ₅ -	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	337	n-Bu	-	(CH ₂) ₅ -	O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, H
	338	n-Bu	-	(CH ₂) ₅ -	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
15	339	n-Bu	-	(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (n-Bu)	H, H
	340	n-Bu	-	(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (i-Bu)	H, H
	341	n-Bu	-	(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	342	n-Bu	-	(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (i-C ₅ H ₁₁)	H, H
	343	n-Bu	-	(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (cy-C ₃ H ₅)	H, H
20	344	n-Bu	-	(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	345	n-Bu	-	(CH ₂) ₅ -	O	-SO ₂ NHCOC ₆ H ₅	CH ₃ , H
	346	n-Bu	-	(CH ₂) ₅ -	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	CH ₃ , H
	347	n-Bu	-	(CH ₂) ₅ -	O	-SO ₂ NHCO (cy-C ₃ H ₅)	CH ₃ , H
	348	n-Bu	-	(CH ₂) ₅ -	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	CH ₃ , H
25	349	n-Bu	-	(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (n-Bu)	CH ₃ , H
	350	n-Bu	-	(CH ₂) ₅ -	O	-SO ₂ NHCONH (n-Bu)	CH ₃ , H
	351	n-Bu	-	(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (i-Bu)	CH ₃ , H
	352	n-Bu	-	(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (n-C ₅ H ₁₁)	CH ₃ , H
	353	n-Bu	-	(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (i-C ₅ H ₁₁)	CH ₃ , H
30	354	n-Bu	-	(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (cy-C ₃ H ₅)	CH ₃ , H
	355	n-Bu	-	(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCOCH ₂ C ₆ H ₅	CH ₃ , H
	356	n-Bu	-	(CH ₂) ₅ -	O	-SO ₂ NHCOC ₆ H ₅	Cl, H
	357	n-Bu	-	(CH ₂) ₅ -	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	Cl, H
	358	n-Bu	-	(CH ₂) ₅ -	O	-SO ₂ NHCO (cy-C ₃ H ₅)	Cl, H

EX.	R ⁶	R ⁷	R ⁸	R ⁹ R ¹⁰	R ¹³	R ² , R ³
	359	n-Bu	-(CH ₂) ₅ -	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	Cl, H
	360	n-Bu	-(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (n-Bu)	Cl, H
	361	n-Bu	-(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (i-Bu)	Cl, H
5	362	n-Bu	-(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (n-C ₅ H ₁₁)	Cl, H
	363	n-Bu	-(CH ₂) ₅ -	O	-SO ₂ NHCONH (n-C ₅ H ₁₁)	Cl, H
	364	n-Bu	-(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (i-C ₅ H ₁₁)	Cl, H
	365	n-Bu	-(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (cy-C ₃ H ₅)	Cl, H
	366	n-Bu	-(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCOCH ₂ C ₆ H ₅	Cl, H
10	367	n-Bu	-(CH ₂) ₅ -	O	-SO ₂ NHCOC ₆ H ₅	F, H
	368	n-Bu	-(CH ₂) ₅ -	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	F, H
	369	n-Bu	-(CH ₂) ₅ -	O	-SO ₂ NHCO (cy-C ₃ H ₅)	F, H
	370	n-Bu	-(CH ₂) ₅ -	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	F, H
	371	n-Bu	-(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (n-Bu)	F, H
15	372	n-Bu	-(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (i-Bu)	F, H
	373	n-Bu	-(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (n-C ₅ H ₁₁)	F, H
	374	n-Bu	-(CH ₂) ₅ -	O	-SO ₂ NHCONH (n-C ₅ H ₁₁)	F, H
	375	n-Bu	-(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (i-C ₅ H ₁₁)	F, H
	376	n-Bu	-(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (cy-C ₃ H ₅)	F, H
20	377	n-Bu	-(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCOCH ₂ C ₆ H ₅	F, H
	378	n-Bu	-(CH ₂) ₅ -	O	-SO ₂ NHCOC ₆ H ₅	H, n-Pr
	379	n-Bu	-(CH ₂) ₅ -	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, n-Pr
	380	n-Bu	-(CH ₂) ₅ -	O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, n-Pr
	381	n-Bu	-(CH ₂) ₅ -	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, n-Pr
25	382	n-Bu	-(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (n-Bu)	H, n-Pr
	383	n-Bu	-(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (i-Bu)	H, n-Pr
	384	n-Bu	-(CH ₂) ₅ -	O	-SO ₂ NHCONH (n-C ₅ H ₁₁)	H, n-Pr
	385	n-Bu	-(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (n-C ₅ H ₁₁)	H, n-Pr
	386	n-Bu	-(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (i-C ₅ H ₁₁)	H, n-Pr
30	387	n-Bu	-(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCO (cy-C ₃ H ₅)	H, n-Pr
	388	n-Bu	-(CH ₂) ₅ -	O	-NH ₂ SO ₂ NHCOCH ₂ C ₆ H ₅	H, n-Pr
	389	n-Bu	-(CH ₂) ₅ -	O	-SO ₂ NHCOC ₆ H ₅	Cl, n-Pr
	390	n-Bu	-(CH ₂) ₅ -	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	Cl, n-Pr
	391	n-Bu	-(CH ₂) ₅ -	O	-SO ₂ NHCO (cy-C ₃ H ₅)	F, n-Pr

	EX.	R ⁶	R ⁷	R ⁸	R ⁹ R ¹⁰	R ¹³	R ² , R ³
	392	n-Bu	-(CH ₂) ₅ -		O	-SO ₂ NHCOCH ₂ C ₆ H ₅	F, n-Pr
	393	n-Bu	-(CH ₂) ₅ -		O	-NHSO ₂ NHCO (n-Bu)	Cl, n-Pr
	394	n-Bu	-(CH ₂) ₅ -		O	-SO ₂ NHCONH (n-Bu)	Cl, n-Pr
5	395	n-Bu	-(CH ₂) ₅ -		O	-SO ₂ NHCONH (i-Bu)	Cl, n-Pr
	396	n-Bu	-(CH ₂) ₅ -		O	-NHSO ₂ NHCO (i-Bu)	F, n-Pr
	397	n-Bu	-(CH ₂) ₅ -		O	-NHSO ₂ NHCO (n-C ₅ H ₁₁)	Cl, n-Pr
	398	n-Bu	-(CH ₂) ₅ -		O	-NHSO ₂ NHCO (i-C ₅ H ₁₁)	F, n-Pr
	399	n-Bu	-(CH ₂) ₅ -		O	-NHSO ₂ NHCO (cy-C ₃ H ₅)	F, n-Pr
10	400	n-Bu	-(CH ₂) ₅ -		O	-NHSO ₂ NHCOCH ₂ C ₆ H ₅	Cl, n-Pr
	401	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-CONHSO ₂ CH ₂ C ₆ H ₅	H, H
	402	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-SO ₂ NHCOC ₆ H ₅	H, H
	403	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	404	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, H
15	405	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	406	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-NHSO ₂ NHCO (n-Bu)	H, H
	407	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-NHSO ₂ NHCO (i-Bu)	H, H
	408	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-NHSO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	409	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-NHSO ₂ NHCO (i-C ₅ H ₁₁)	H, H
20	410	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-NHSO ₂ NHCO (cy-C ₃ H ₅)	H, H
	411	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-NHSO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	412	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-SO ₂ NHCOC ₆ H ₅	CH ₃ , H
	413	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-SO ₂ NHCO (n-C ₅ H ₁₁)	CH ₃ , H
	414	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-SO ₂ NHCO (cy-C ₃ H ₅)	CH ₃ , H
25	415	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-SO ₂ NHCOCH ₂ C ₆ H ₅	CH ₃ , H
	416	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-NHSO ₂ NHCO (n-Bu)	CH ₃ , H
	417	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-SO ₂ NHCONH (n-Bu)	CH ₃ , H
	418	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-NHSO ₂ NHCO (i-Bu)	CH ₃ , H
	419	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-NHSO ₂ NHCO (n-C ₅ H ₁₁)	CH ₃ , H
30	420	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-NHSO ₂ NHCO (i-C ₅ H ₁₁)	CH ₃ , H
	421	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-NHSO ₂ NHCO (cy-C ₃ H ₅)	CH ₃ , H
	422	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-NHSO ₂ NHCOCH ₂ C ₆ H ₅	CH ₃ , H
	423	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-SO ₂ NHCOC ₆ H ₅	Cl, H
	424	n-Pr	-(CH ₂) S (CH ₂) ₂ -		O	-SO ₂ NHCO (n-C ₅ H ₁₁)	Cl, H

	EX.	R ⁶	R ⁷	R ⁸	R ⁹ R ¹⁰	R ¹³	R ² , R ³	
	425	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-SO ₂ NHCO (cy-C ₃ H ₅)	Cl, H	
	426	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	Cl, H	a
	427	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-NHSO ₂ NHCO (n-Bu)	Cl, H	
5	428	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-NHSO ₂ NHCO (n-Bu)	Cl, H	*
	429	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-SO ₂ NHCONH (i-Bu)	Cl, H	
	430	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-NHSO ₂ NHCO (n-C ₅ H ₁₁)	Cl, H	
	431	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-NHSO ₂ NHCO (i-C ₅ H ₁₁)	Cl, H	
	432	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-NHSO ₂ NHCO (cy-C ₃ H ₅)	Cl, H	
10	433	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-NHSO ₂ NHCOCH ₂ C ₆ H ₅	Cl, H	
	434	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-SO ₂ NHCOC ₆ H ₅	F, H	
	435	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	F, H	
	436	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-SO ₂ NHCO (cy-C ₃ H ₅)	F, H	
	437	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	F, H	
15	438	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-NHSO ₂ NHCO (n-Bu)	F, H	
	439	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-SO ₂ NHCONH (n-Bu)	F, H	
	440	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-NHSO ₂ NHCO (i-Bu)	F, H	
	441	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-NHSO ₂ NHCO (n-C ₅ H ₁₁)	F, H	
	442	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-NHSO ₂ NHCO (i-C ₅ H ₁₁)	F, H	
20	443	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-NHSO ₂ NHCO (cy-C ₃ H ₅)	F, H	
	444	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-NHSO ₂ NHCOCH ₂ C ₆ H ₅	F, H	
	445	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-SO ₂ NHCOC ₆ H ₅	H, n-Pr	
	446	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, n-Pr	
	447	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, n-Pr	
25	448	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, n-Pr	
	449	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-NHSO ₂ NHCO (n-Bu)	H, n-Pr	
	450	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-SO ₂ NHCONH (n-Bu)	H, n-Pr	
	451	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-NHSO ₂ NHCO (i-Bu)	H, n-Pr	
	452	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-NHSO ₂ NHCO (n-C ₅ H ₁₁)	H, n-Pr	
30	453	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-NHSO ₂ NHCO (i-C ₅ H ₁₁)	H, n-Pr	
	454	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-NHSO ₂ NHCO (cy-C ₃ H ₅)	H, n-Pr	
	455	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-NHSO ₂ NHCOCH ₂ C ₆ H ₅	H, n-Pr	b
	456	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-SO ₂ NHCOC ₆ H ₅	Cl, n-Pr	
	457	n-Pr	-(CH ₂)	S (CH ₂) ₂ -	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	Cl, n-Pr	c

	EX.	R ⁶	R ⁷	R ⁸	R ⁹ R ¹⁰	R ¹³	R ² , R ³
	458	n-Pr	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-SO ₂ NHCO (cy-C ₃ H ₅)	F, n-Pr
	459	n-Pr	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	F, n-Pr
	460	n-Pr	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-NHSO ₂ NHCO (n-Bu)	Cl, n-Pr
5	461	n-Pr	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-SO ₂ NHCONH (n-Bu)	Cl, n-Pr
	462	n-Pr	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-NHSO ₂ NHCO (i-Bu)	F, n-Pr
	463	n-Pr	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-NHSO ₂ NHCO (n-C ₅ H ₁₁)	Cl, n-Pr
	463	n-Pr	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-NHSO ₂ NHCO (i-C ₅ H ₁₁)	F, n-Pr
	464	n-Pr	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-NHSO ₂ NHCO (cy-C ₃ H ₅)	F, n-Pr
10	465	n-Pr	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-NHSO ₂ NHCOCH ₂ C ₆ H ₅	Cl, n-Pr
	466	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-SO ₂ NHCOC ₆ H ₅	H, H
	467	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	468	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, H
	469	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
15	470	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-NHSO ₂ NHCO (n-Bu)	H, H
	471	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-NHSO ₂ NHCO (i-Bu)	H, H
	472	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-NHSO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	473	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-NHSO ₂ NHCO (i-C ₅ H ₁₁)	H, H
	474	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-NHSO ₂ NHCO (cy-C ₃ H ₅)	H, H
20	475	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-NHSO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	476	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-SO ₂ NHCOC ₆ H ₅	CH ₃ , H
	477	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	CH ₃ , H
	478	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-SO ₂ NHCO (cy-C ₃ H ₅)	CH ₃ , H
	479	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	CH ₃ , H
25	480	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-NHSO ₂ NHCO (n-Bu)	CH ₃ , H
	481	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-SO ₂ NHCONH (n-Bu)	CH ₃ , H
	482	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-NHSO ₂ NHCO (i-Bu)	CH ₃ , H
	483	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-NHSO ₂ NHCO (n-C ₅ H ₁₁)	CH ₃ , H
	484	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-NHSO ₂ NHCO (i-C ₅ H ₁₁)	CH ₃ , H
30	485	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-NHSO ₂ NHCO (cy-C ₃ H ₅)	CH ₃ , H
	486	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-NHSO ₂ NHCOCH ₂ C ₆ H ₅	CH ₃ , H
	487	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-SO ₂ NHCOC ₆ H ₅	Cl, H
	488	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	Cl, H
	489	n-Bu	-(CH ₂) ₂ S	-(CH ₂) ₂	O	-SO ₂ NHCO (cy-C ₃ H ₅)	Cl, H

	EX.	R ⁶	R ⁷	R ⁸	R ⁹ R ¹⁰	R ¹³	R ² , R ³
	490	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	Cl, H
	491	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-NH ₂ SO ₂ NHCO (n-Bu)	Cl, H
	492	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-NH ₂ SO ₂ NHCO (i-Bu)	Cl, H
5	493	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-NH ₂ SO ₂ NHCO (n-C ₅ H ₁₁)	Cl, H
	494	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-SO ₂ NHCONH (n-C ₅ H ₁₁)	Cl, H
	495	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-NH ₂ SO ₂ NHCO (i-C ₅ H ₁₁)	Cl, H
	496	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-NH ₂ SO ₂ NHCO (cy-C ₃ H ₅)	Cl, H
	497	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-NH ₂ SO ₂ NHCOCH ₂ C ₆ H ₅	Cl, H
10	498	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-SO ₂ NHCOC ₆ H ₅	F, H
	499	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	F, H
	500	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-SO ₂ NHCO (cy-C ₃ H ₅)	F, H
	501	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	F, H
	502	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-NH ₂ SO ₂ NHCO (n-Bu)	F, H
15	503	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-NH ₂ SO ₂ NHCO (i-Bu)	F, H
	504	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-NH ₂ SO ₂ NHCO (n-C ₅ H ₁₁)	F, H
	505	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-SO ₂ NHCONH (n-C ₅ H ₁₁)	F, H
	506	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-NH ₂ SO ₂ NHCO (i-C ₅ H ₁₁)	F, H
	507	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-NH ₂ SO ₂ NHCO (cy-C ₃ H ₅)	F, H
20	508	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-NH ₂ SO ₂ NHCOCH ₂ C ₆ H ₅	F, H
	509	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-SO ₂ NHCOC ₆ H ₅	H, n-Pr
	510	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, n-Pr
	511	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, n-Pr
	512	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, n-Pr
25	513	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-NH ₂ SO ₂ NHCO (n-Bu)	H, n-Pr
	514	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-NH ₂ SO ₂ NHCO (i-Bu)	H, n-Pr
	515	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-SO ₂ NHCONH (n-C ₅ H ₁₁)	H, n-Pr
	516	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-NH ₂ SO ₂ NHCO (n-C ₅ H ₁₁)	H, n-Pr
	517	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-NH ₂ SO ₂ NHCO (i-C ₅ H ₁₁)	H, n-Pr
30	518	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-NH ₂ SO ₂ NHCO (cy-C ₃ H ₅)	H, n-Pr
	519	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-NH ₂ SO ₂ NHCOCH ₂ C ₆ H ₅	H, n-Pr
	520	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-SO ₂ NHCOC ₆ H ₅	Cl, n-Pr
	521	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	Cl, n-Pr
	522	n-Bu	-	(CH ₂) S (CH ₂) 2-	O	-SO ₂ NHCO (cy-C ₃ H ₅)	F, n-Pr

	EX.	R ⁶	R ⁷	R ⁸	R ⁹ R ¹⁰	R ¹³	R ² , R ³
	523	n-Bu	-(CH ₂)	S(CH ₂) ₂ -	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	F, n-Pr
	524	n-Bu	-(CH ₂)	S(CH ₂) ₂ -	O	-NHSO ₂ NHCO (n-Bu)	Cl, n-Pr
	525	n-Bu	-(CH ₂)	S(CH ₂) ₂ -	O	-SO ₂ NHCONH (n-Bu)	Cl, n-Pr
5	526	n-Bu	-(CH ₂)	S(CH ₂) ₂ -	O	-SO ₂ NHCONH (i-Bu)	Cl, n-Pr
	527	n-Bu	-(CH ₂)	S(CH ₂) ₂ -	O	-NHSO ₂ NHCO (i-Bu)	F, n-Pr
	528	n-Bu	-(CH ₂)	S(CH ₂) ₂ -	O	-NHSO ₂ NHCO (n-C ₅ H ₁₁)	Cl, n-Pr
	529	n-Bu	-(CH ₂)	S(CH ₂) ₂ -	O	-NHSO ₂ NHCO (i-C ₅ H ₁₁)	F, n-Pr
	530	n-Bu	-(CH ₂)	S(CH ₂) ₂ -	O	-NHSO ₂ NHCO (cy-C ₃ H ₅)	F, n-Pr
10	531	n-Bu	-(CH ₂)	S(CH ₂) ₂ -	O	-NHSO ₂ NHCOCH ₂ C ₆ H ₅	Cl, n-Pr
	532	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-CONHSO ₂ CH ₂ C ₆ H ₅	H, H
	533	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-SO ₂ NHCOC ₆ H ₅	H, H
	534	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	535	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, H
15	536	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	537	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-NHSO ₂ NHCO (n-Bu)	H, H
	538	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-NHSO ₂ NHCO (i-Bu)	H, H
	539	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-NHSO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	540	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-NHSO ₂ NHCO (i-C ₅ H ₁₁)	H, H
20	541	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-NHSO ₂ NHCO (cy-C ₃ H ₅)	H, H
	542	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-NHSO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	543	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-CONHSO ₂ CH ₂ C ₆ H ₅	H, H
	544	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-SO ₂ NHCOC ₆ H ₅	H, H
	545	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, H
25	546	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, H
	547	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	548	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-NHSO ₂ NHCO (n-Bu)	H, H
	549	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-NHSO ₂ NHCO (i-Bu)	H, H
	550	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-NHSO ₂ NHCO (n-C ₅ H ₁₁)	H, H
30	551	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-NHSO ₂ NHCO (i-C ₅ H ₁₁)	H, H
	552	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-NHSO ₂ NHCO (cy-C ₃ H ₅)	H, H
	553	n-Pr	-(CH ₂)	O(CH ₂) ₂ -	O	-NHSO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	554	n-Pr	-(CH ₂)	SO ₂ (CH ₂) ₂ -	O	-CONHSO ₂ CH ₂ C ₆ H ₅	H, H
	555	n-Pr	-(CH ₂)	SO ₂ (CH ₂) ₂ -	O	-SO ₂ NHCOC ₆ H ₅	H, H

	EX.	R ⁶	R ⁷	R ⁸	R ⁹	R ¹⁰	R ¹³	R ² , R ³
	556	n-Pr	-	(CH ₂) SO ₂ (CH ₂) ₂ -	O		-SO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	557	n-Pr	-	(CH ₂) SO ₂ (CH ₂) ₂ -	O		-SO ₂ NHCO (cy-C ₃ H ₅)	H, H
	558	n-Pr	-	(CH ₂) SO ₂ (CH ₂) ₂ -	O		-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
5	559	n-Pr	-	(CH ₂) SO ₂ (CH ₂) ₂ -	O		-NHSO ₂ NHCO (n-Bu)	H, H
	560	n-Pr	-	(CH ₂) SO ₂ (CH ₂) ₂ -	O		-NHSO ₂ NHCO (i-Bu)	H, H
	561	n-Pr	-	(CH ₂) SO ₂ (CH ₂) ₂ -	O		-NHSO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	562	n-Pr	-	(CH ₂) SO ₂ (CH ₂) ₂ -	O		-NHSO ₂ NHCO (i-C ₅ H ₁₁)	H, H
	563	n-Pr	-	(CH ₂) SO ₂ (CH ₂) ₂ -	O		-NHSO ₂ NHCO (cy-C ₃ H ₅)	H, H
10	564	n-Pr	-	(CH ₂) SO ₂ (CH ₂) ₂ -	O		-NHSO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	565	n-Pr	-	(CH ₂) NCOMe (CH ₂) ₂ -	O		-CONHSO ₂ CH ₂ C ₆ H ₅	H, H
	566	n-Pr	-	(CH ₂) NCOMe (CH ₂) ₂ -	O		-SO ₂ NHCOC ₆ H ₅	H, H
	567	n-Pr	-	(CH ₂) NCOMe (CH ₂) ₂ -	O		-SO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	568	n-Pr	-	(CH ₂) NCOMe (CH ₂) ₂ -	O		-SO ₂ NHCO (cy-C ₃ H ₅)	H, H
15	569	n-Pr	-	(CH ₂) NCOMe (CH ₂) ₂ -	O		-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	570	n-Pr	-	(CH ₂) NCOMe (CH ₂) ₂ -	O		-NHSO ₂ NHCO (n-Bu)	H, H
	571	n-Pr	-	(CH ₂) NCOMe (CH ₂) ₂ -	O		-NHSO ₂ NHCO (i-Bu)	H, H
	572	n-Pr	-	(CH ₂) NCOMe (CH ₂) ₂ -	O		-NHSO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	573	n-Pr	-	(CH ₂) NCOMe (CH ₂) ₂ -	O		-NHSO ₂ NHCO (i-C ₅ H ₁₁)	H, H
20	574	n-Pr	-	(CH ₂) NCOMe (CH ₂) ₂ -	O		-NHSO ₂ NHCO (cy-C ₃ H ₅)	H, H
	575	n-Pr	-	(CH ₂) NCOMe (CH ₂) ₂ -	O		-NHSO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	576	n-Pr	-	(CH ₂) NCOPh (CH ₂) ₂ -	O		-CONHSO ₂ CH ₂ C ₆ H ₅	H, H
	577	n-Pr	-	(CH ₂) NCOPh (CH ₂) ₂ -	O		-SO ₂ NHCOC ₆ H ₅	H, H
	578	n-Pr	-	(CH ₂) NCOPh (CH ₂) ₂ -	O		-SO ₂ NHCO (n-C ₅ H ₁₁)	H, H
25	579	n-Pr	-	(CH ₂) NCOPh (CH ₂) ₂ -	O		-SO ₂ NHCO (cy-C ₃ H ₅)	H, H
	580	n-Pr	-	(CH ₂) NCOPh (CH ₂) ₂ -	O		-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	581	n-Pr	-	(CH ₂) NCOPh (CH ₂) ₂ -	O		-NHSO ₂ NHCO (n-Bu)	H, H
	582	n-Pr	-	(CH ₂) NCOPh (CH ₂) ₂ -	O		-NHSO ₂ NHCO (i-Bu)	H, H
	583	n-Pr	-	(CH ₂) NCOPh (CH ₂) ₂ -	O		-NHSO ₂ NHCO (n-C ₅ H ₁₁)	H, H
30	584	n-Pr	-	(CH ₂) NCOPh (CH ₂) ₂ -	O		-NHSO ₂ NHCO (i-C ₅ H ₁₁)	H, H
	585	n-Pr	-	(CH ₂) NCOPh (CH ₂) ₂ -	O		-NHSO ₂ NHCO (cy-C ₃ H ₅)	H, H
	586	n-Pr	-	(CH ₂) NCOPh (CH ₂) ₂ -	O		-NHSO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	587	n-Pr	-	(CH ₂) NCH ₂ Ph (CH ₂) ₂ -	O		-CONHSO ₂ CH ₂ C ₆ H ₅	H, H
	588	n-Pr	-	(CH ₂) NCH ₂ Ph (CH ₂) ₂ -	O		-SO ₂ NHCOC ₆ H ₅	H, H

	EX.	R ⁶	R ⁷	R ⁸	R ⁹ R ¹⁰	R ¹³	R ² , R ³
	589	n-Pr	-(CH ₂)	NCH ₂ Ph (CH ₂) ₂ -	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	590	n-Pr	-(CH ₂)	NCH ₂ Ph (CH ₂) ₂ -	O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, H
	591	n-Pr	-(CH ₂)	NCH ₂ Ph (CH ₂) ₂ -	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
5	592	n-Pr	-(CH ₂)	NCH ₂ Ph (CH ₂) ₂ -	O	-NHSO ₂ NHCO (n-Bu)	H, H
	593	n-Pr	-(CH ₂)	NCH ₂ Ph (CH ₂) ₂ -	O	-NHSO ₂ NHCO (i-Bu)	H, H
	594	n-Pr	-(CH ₂)	NCH ₂ Ph (CH ₂) ₂ -	O	-NHSO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	595	n-Pr	-(CH ₂)	NCH ₂ Ph (CH ₂) ₂ -	O	-NHSO ₂ NHCO (i-C ₅ H ₁₁)	H, H
	596	n-Pr	-(CH ₂)	NCH ₂ Ph (CH ₂) ₂ -	O	-NHSO ₂ NHCO (cy-C ₃ H ₅)	H, H
10	597	n-Pr	-(CH ₂)	NCH ₂ Ph (CH ₂) ₂ -	O	-NHSO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	598	n-Pr	CF ₃	CF ₃	O	-SO ₂ NHCOC ₆ H ₅	H, H
	599	n-Pr	CF ₃	CF ₃	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	600	n-Pr	CF ₃	CF ₃	O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, H
	601	n-Pr	CF ₃	CF ₃	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
15	602	n-Pr	CF ₃	CF ₃	O	-NHSO ₂ NHCO (n-Bu)	H, H
	603	n-Pr	CF ₃	CF ₃	O	-NHSO ₂ NHCO (i-Bu)	H, H
	604	n-Pr	CF ₃	CF ₃	O	-NHSO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	605	n-Pr	CF ₃	CF ₃	O	-NHSO ₂ NHCO (i-C ₅ H ₁₁)	H, H
	606	n-Pr	CF ₃	CF ₃	O	-NHSO ₂ NHCO (cy-C ₃ H ₅)	H, H
20	607	n-Pr	CF ₃	CF ₃	O	-NHSO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	608	n-Pr	Ph	Ph	O	-SO ₂ NHCOC ₆ H ₅	H, H
	609	n-Pr	Ph	Ph	O	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	610	n-Pr	Ph	Ph	O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, H
	611	n-Pr	Ph	Ph	O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
25	612	n-Pr	Ph	Ph	O	-NHSO ₂ NHCO (n-Bu)	H, H
	613	n-Pr	Ph	Ph	O	-NHSO ₂ NHCO (i-Bu)	H, H
	614	n-Pr	Ph	Ph	O	-NHSO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	615	n-Pr	Ph	Ph	O	-NHSO ₂ NHCO (i-C ₅ H ₁₁)	H, H
	616	n-Pr	Ph	Ph	O	-NHSO ₂ NHCO (cy-C ₃ H ₅)	H, H
30	617	n-Pr	Ph	Ph	O	-NHSO ₂ NHCOCH ₂ C ₆ H ₅	H, H
	618	n-Pr	-(CH ₂) ₂ -		O	-SO ₂ NHCOC ₆ H ₅	H, H
	619	n-Pr	-(CH ₂) ₂ -		O	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, H
	620	n-Pr	-(CH ₂) ₂ -		O	-SO ₂ NHCO (cy-C ₃ H ₅)	H, H
	621	n-Pr	-(CH ₂) ₂ -		O	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H

EX.	R ⁶	R ⁷	R ⁸	R ⁹ R ¹⁰	R ¹³	R ² , R ³
622	n-Pr	-(CH ₂) ₂ -		O	-NHSO ₂ NHCO (n-Bu)	H, H
623	n-Pr	-(CH ₂) ₂ -		O	-NHSO ₂ NHCO (i-Bu)	H, H
624	n-Pr	-(CH ₂) ₂ -		O	-NHSO ₂ NHCO (n-C ₅ H ₁₁)	H, H
5 625	n-Pr	-(CH ₂) ₂ -		O	-NHSO ₂ NHCO (i-C ₅ H ₁₁)	H, H
626	n-Pr	-(CH ₂) ₂ -		O	-NHSO ₂ NHCO (cy-C ₃ H ₅)	H, H
627	n-Pr	-(CH ₂) ₂ -		O	-NHSO ₂ NHCOCH ₂ C ₆ H ₅	H, H
628	n-Pr	CH ₃	CH ₃	S	-SO ₂ NHCOCH ₂ C ₆ H ₅	H, H
629	n-Pr	CH ₃	CH ₃	S	-CONHSO ₂ CH ₂ C ₆ H ₅	H, H
10 630	n-Pr	CH ₃	CH ₃	S	-SO ₂ NHCO (n-C ₅ H ₁₁)	H, H

Utility

Angiotensin II (AII) produces numerous biological responses (e.g., vasoconstriction) through stimulation of its receptors on cell membranes. For the purpose of identifying compounds such as AII antagonists which are capable of interacting with the AII receptor, a ligand-receptor binding assay was utilized for the initial screen. The assay was carried out according to the method described by Chiu, et al., Receptor, 1 33, (1990). In brief, aliquots of a freshly prepared particulate fraction of rat adrenal cortex were incubated with 0.05 nM [¹²⁵I] AII and varying concentrations of potential AII antagonists in a Tris buffer. After a 1 h incubation the reaction was terminated by addition of cold assay buffer. The bound and free radioactivity were rapidly separated through glass-fiber filters, and the trapped radioactivity was quantitated by scintillation counting. The inhibitory concentration (IC₅₀) of potential AII antagonists which gives 50% displacement of the total specifically bound [¹²⁵I] AII is presented as a measure of the affinity of such compound for the AII receptor.

Using the assay method described above, the compounds of this invention are found to exhibit an

activity of at least $IC_{50} < 10$ micromolar, thereby demonstrating and confirming the activity of these compounds as effective AII antagonists.

The potential antihypertensive effects of the compounds of this invention may be demonstrated by administering the compounds to awake rats made hypertensive by ligation of the left renal artery [Cangiano et al., *J. Pharmacol. Exp. Ther.*, **1979**, 208, 310]. This procedure increases blood pressure by increasing renin production with consequent elevation of AII levels. Compounds are administered intravenously via a cannula in the jugular vein at 10 mg/kg. Arterial blood pressure is continuously measured directly through a carotid artery cannula and recorded using a pressure transducer and a polygraph. Blood pressure levels after treatment are compared to pretreatment levels to determine the antihypertensive effects of the compounds.

Using the in vivo methodology described above, the compounds of this invention are found to exhibit an activity (intravenous) which is 10 mg/kg or less, and/or an activity (oral) which is 100 mg/kg or less, thereby demonstrating and confirming the utility of these compounds as effective agents in lowering blood pressure.

The compounds of the invention are useful in treating hypertension. They are also of value in the management of acute and chronic congestive heart failure and angina. These compounds may also be expected to be useful in the treatment of primary and secondary hyperaldosteronism; renal diseases such as diabetic nephropathy, glomerulonephritis, glomerular sclerosis, nephrotic syndrome, hypertensive nephrosclerosis, end stage renal disease, used in renal transplant therapy, and to treat renovascular hypertension, scleroderma,

left ventricular dysfunction, systolic and diastolic dysfunction, diabetic retinopathy and in the management of vascular disorders such as migraine, Raynaud's disease, and as prophylaxis to minimize the

5 atherosclerotic process and neointimal hyperplasia following angioplasty or vascular injury and to retard the onset of type II diabetes. The application of the compounds of this invention for these and similar disorders will be apparent to those skilled in the art.

10 The compounds of this invention are also useful to treat elevated intraocular pressure and to enhance retinal blood flow and can be administered to patients in need of such treatment with typical pharmaceutical formulations such as tablets, capsules, injectables and

15 the like as well as topical ocular formulations in the form of solutions, ointments, inserts, gels and the like. Pharmaceutical formulations prepared to treat intraocular pressure would typically contain about 0.1% to 15% by weight, preferably 0.5% to 2% by weight, of a

20 compound of this invention. For this use, the compounds of this invention may also be used in combination with other medications for the treatment of glaucoma including choline esterase inhibitors such as physostigmine salicylate or demecarium bromide,

25 parasympathomimetic agents such as pilocarpine nitrate, β -adrenergic antagonists such as timolol maleate, adrenergic agonists such as epinephrine and carbonic anhydrase inhibitors such as MK-507.

In the management of hypertension and the clinical

30 conditions noted above, the compounds of this invention may be utilized with a pharmaceutical carrier in compositions such as tablets, capsules or elixirs for oral administration, suppositories for rectal administration, sterile solutions or suspensions for

parenteral or intramuscular administration, and the like. The compounds of this invention can be administered to patients (animals and human) in need of such treatment in dosages that will provide optimal pharmaceutical efficacy. Although the dose will vary from patient to patient depending upon the nature and severity of disease, the patient's weight, special diet that is being followed by a patient, concurrent medication, and other factors which those skilled in the art will recognize, the dosage range will generally be about 1 to 1000 mg per patient per day which can be administered in single or multiple doses. Preferably, the dosage range will be about 5 to 500 mg per patient per day; more preferably about 5 to 300 mg per patient per day.

The compounds of this invention can also be administered in combination with other antihypertensives and/or diuretics. For example, the compounds of this invention can be given in combination with diuretics such as hydrochlorothiazide, chlorothiazide, chlorthalidone, methylclothiazide, furosemide, ethacrynic acid, triamterene, amiloride spironolactone and atriopeptin; calcium channel blockers, such as diltiazem, felodipine, nifedipine, amlodipine, nimodipine, isradipine, nitrendipine and verapamil; β -adrenergic antagonists such as timolol, atenolol, metoprolol, propranolol, nadolol and pindolol; angiotensin converting enzyme inhibitors such as enalapril, lisinopril, captopril, ramipril, quinapril and zofenopril; renin inhibitors such as A-69729, FK 906 and FK 744; α -adrenergic antagonists such as prazosin, doxazosin, and terazosin; sympatholytic agents such as methyldopa, clonidine and guanabenz; atriopeptidase inhibitors (alone or with ANP) such as UK-79300;

serotonin antagonists such as ketanserin; A₂-adrenosine receptor agonists such as CGS 22492C; potassium channel agonists such as pinacidil and cromakalim; and various other antihypertensive drugs including reserpine, minoxidil, guanethidine, hydralazine hydrochloride and sodium nitroprusside as well as combinations of the above-named drugs. Combinations useful in the management of congestive heart failure include, in addition, compounds of this invention with cardiac stimulants such as dobutamine and xamoterol and phosphodiesterase inhibitors including amrinone and milrinone.

Typically, the individual daily dosages for these combinations can range from about one-fifth of the minimally recommended clinical dosages to the maximum recommended levels for the entities when they are given singly. To illustrate these combinations, one of the angiotensin II antagonists of this invention effective clinically in the 5-500 milligrams per day range can be effectively combined at levels at the 1.0-500 milligrams per day range with the following compounds at the indicated per day dose range; hydrochlorothiazide (6-100 mg), chlorothiazide (125-500 mg), ethacrynic acid (5-200 mg), amiloride (5-20 mg), furosemide (5-80 mg), propranolol (10-480 mg), timolol maleate (1-20 mg), methyldopa (125-2000 mg), felodipine (1-20 mg), nifedipine (5-120 mg), nitrendipine (5-60 mg), and diltiazem (30-540 mg). In addition, triple drug combinations of hydrochlorothiazide (5-100 mg) plus amiloride (5-20 mg) plus angiotensin II antagonists of this invention (1-500 mg) or hydrochlorothiazide (5-100 mg) plus timolol maleate (5-60 mg) plus an angiotensin II antagonists of this invention (1-500 mg) or hydrochlorothiazide (5-200 mg) and nifedipine (5-60 mg)

plus an angiotensin II antagonist of this invention (1-500 mg) are effective combinations to control blood pressure in hypertensive patients. Naturally, these dose ranges can be adjusted on a unit basis as necessary to permit divided daily dosage and, as noted above, the dose will vary depending on the nature and severity of the disease, weight of patient, special diets and other factors.

The active ingredient can be administered orally in solid dosage forms, such as capsules, tablets, and powders, or in liquid dosage forms, such as elixirs, syrups, and suspensions. It can also be administered parenterally, in sterile liquid dosage forms.

Gelatin capsules contain the active ingredient and powdered carriers, such as lactose, starch, cellulose derivatives, magnesium stearate, stearic acid, and the like. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of hours. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the tablet from the atmosphere, or enteric coated for selective disintegration in the gastrointestinal tract.

Liquid dosage forms for oral administration can contain coloring and flavoring to increase patient acceptance.

In general, water, a suitable oil, saline, aqueous dextrose (glucose), and related sugar solutions and glycols such as propylene glycol or polyethylene glycols are suitable carriers for parenteral solutions. Solutions for parenteral administration preferably contain a water soluble salt of the active ingredient, suitable stabilizing agents, and if necessary, buffer

substances. Antioxidizing agents such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined, are suitable stabilizing agents. Also used are citric acid and its salts and sodium EDTA.

5 In addition, parenteral solutions can contain preservatives, such as benzalkonium chloride, methyl- or propylparaben, and chlorobutanol.

Suitable pharmaceutical carriers are described in Remington's Pharmaceutical Sciences, A. Osol, a standard
10 reference text in this field.

Useful pharmaceutical dosage-forms for administration of the compounds of this invention can be illustrated as follows:

15 Capsules

A large number of unit capsules are prepared by filling standard two-piece hard gelatin capsules each with 100 milligrams of powdered active ingredient, 150 milligrams of lactose, 50 milligrams of cellulose, and 6
20 milligrams magnesium stearate.

Soft Gelatin Capsules

A mixture of active ingredient in a digestible oil such as soybean oil, cottonseed oil or olive oil is
25 prepared and injected by means of a positive displacement pump into gelatin to form soft gelatin capsules containing 100 milligrams of the active ingredient. The capsules are washed and dried.

30 Tablets

A large number of tablets are prepared by conventional procedures so that the dosage unit is 100 milligrams of active ingredient, 0.2 milligrams of colloidal silicon dioxide, 5 milligrams of magnesium

stearate, 275 milligrams of microcrystalline cellulose, 11 milligrams of starch and 98.8 milligrams of lactose. Appropriate coatings may be applied to increase palatability or delay absorption.

5

Injectable

A parenteral composition suitable for administration by injection is prepared by stirring 1.5% by weight of active ingredient in 10% by volume propylene glycol. The solution is made to volume with water for injection and sterilized.


Suspension

An aqueous suspension is prepared for oral administration so that each 5 milliliters contain 100 milligrams of finely divided active ingredient, 100 milligrams of sodium carboxymethyl cellulose, 5 milligrams of sodium benzoate, 1.0 grams of sorbitol solution, U.S.P., and 0.025 milliliters of vanillin.

The same dosage forms can generally be used when the compounds of this invention are administered stepwise in conjunction with another therapeutic agent. When the drugs are administered in physical combination, the dosage form and administration route should be selected for compatibility with both drugs.


What is claimed is:

1. A compound of formula (I)



5 wherein:

R¹ is other than in the ortho position and is:



R² is

- | | |
|----|--|
| 10 | (a) H, |
| | (b) halo (F, Cl, Br, I), |
| | (c) C ₁ -C ₄ alkyl, |
| | (d) C ₁ -C ₄ alkoxy, |
| | (e) C ₁ -C ₄ acyloxy, |
| 15 | (f) C ₁ -C ₄ alkylthio, |
| | (g) C ₁ -C ₄ alkylsulfinyl, |
| | (h) C ₁ -C ₄ alkylsulfonyl, |
| | (i) hydroxy (C ₁ -C ₄) alkyl, |
| | (j) aryl (C ₁ -C ₄) alkyl, |
| 20 | (k) -CO ₂ H, |
| | (l) -CN, |
| | (m) -CONHOR ¹² , |
| | (n) -SO ₂ NHR ²¹ , |
| | (o) -NH ₂ , |
| 25 | (p) C ₁ -C ₄ alkylamino, |

- (q) C₁-C₄ dialkylamino,
(r) -NHSO₂R²⁰,
(s) -NO₂,
(t) furyl,
5 (u) aryl;

wherein aryl is phenyl optionally substituted with one or two substituents selected from the group consisting of halo, C₁-C₄ alkyl, C₁-C₄ alkoxy, -NO₂, -CF₃, C₁-C₄ alkylthio, -OH, -NH₂, C₁-C₄ alkylamino, C₁-C₄ dialkylamino, -CN, -CO₂H, -CO₂CH₃, -CO₂CH₂CH₃, -CO₂-
10 benzyl;

R³ is

- (a) H,
(b) halo,
15 (c) C₁-C₄ alkyl,
(d) C₁-C₄ alkoxy,
(e) C₁-C₄ alkoxyalkyl;

R⁴ is

- (a) -CN,
20 (b) -NO₂,
(c) -CO₂R¹¹;

R⁵ is

- (a) H,
(b) C₁-C₆ alkyl,
25 (c) C₃-C₆ cycloalkyl,
(d) C₂-C₄ alkenyl,
(e) C₂-C₄ alkynyl;

R⁶ is

- (a) C₁-C₁₀ alkyl,
30 (b) C₃-C₈ alkenyl,
(c) C₃-C₈ alkynyl,
(d) C₃-C₈ cycloalkyl,
(e) C₄-C₈ cycloalkenyl,
(f) C₄-C₁₀ cycloalkylalkyl,

- (g) C₅-C₁₀ cycloalkylalkenyl,
 - (h) C₅-C₁₀ cycloalkylalkynyl,
 - (i) $-(\text{CH}_2)_s\text{Z}(\text{CH}_2)_m\text{R}^5$,
 - (j) phenyl, optionally substituted with 1-2
 5 substituents selected from the group of halo, C₁-C₄
 alkyl, C₁-C₄ alkoxy, -NO₂, -NH₂, -OH and benzyloxy,
 - (k) benzyl, optionally substituted on the phenyl
 ring with 1-2 substituents selected from the group of
 halo, C₁-C₄ alkyl, C₁-C₄ alkoxy and -NO₂;
- 10 R⁷, R⁸, R⁹, and R¹⁰ are independently chosen from
- (a) H,
 - (b) C₁-C₈ alkyl,
 - (c) C₁-C₈ perfluoroalkyl,
 - 15 (d) C₃-C₆ cycloalkyl,
 - (e) -NO₂,
 - (f) -CN,
 - (g) -CONR¹⁵R¹⁶,
 - (h) -CO₂R¹⁷,
 - 20 (i) -OR¹⁸,
 - (j) $-(\text{CH}_2)_n\text{CONR}^{15}\text{R}^{16}$ where n is 1-4,
 - (k) $-(\text{CH}_2)_n\text{CO}_2\text{R}^{17}$ where n is 1-4,
 - (l) $(\text{CH}_2)_n\text{OR}^{18}$ where n is 1-4,
 - (m) aryl, wherein aryl is as defined above,
 - 25 (n) -CH₂ aryl, wherein aryl is as defined above,
 - (o) R⁷ and R⁸ taken together are $-(\text{CH}_2)_t-$, or
 $-(\text{CH}_2)_m\text{X}(\text{CH}_2)_q-$,
 - (p) R⁹ and R¹⁰ taken together can be S, or O;
- R¹¹ is
- 30 (a) H,
 - (b) C₁-C₄ alkyl,
 - (c) C₁-C₄ cycloalkyl,
 - (d) phenyl,
 - (e) benzyl;


R¹² is

- (a) H,
- (b) methyl,
- (c) benzyl;


5 R¹³ is

- (a) -CH₂CO₂H,
- (b) -C(CF₃)₂OH,
- (c) -CONHNHSO₂CF₃,
- (d) -CONHOR¹²,
- 10 (e) -CONHSO₂R²⁰,
- (f) -CONHSO₂NHR¹⁹,
- (g) -C(OH)R¹⁹PO₃H₂,
- (h) -NHCONHSO₂R²⁰,
- (i) -NHPO₃H₂,
- 15 (j) -SO₂NHCOR¹⁹,
- (k) -OPO₃H₂,
- (l) -OSO₃H,
- (m) -PO(OH)R¹⁹,
- (n) -PO₃H₂,
- 20 (o) -SO₃H,
- (p) -SO₂NHR¹⁹,
- (q) -NHSO₂NHCOR¹⁹,
- (r) -SO₂NHCONHR¹⁹,
- (s)


25




(t)



(u)



(v)

R¹⁴ is

- (a) H,
- 5 (b) C₁-C₄ alkyl,
- (c) aryl, wherein aryl is defined as above,
- (d) benzyl,
- (e) COR¹¹,
- (f) CONHR¹¹;

10 R¹⁵ and R¹⁶ are independently

- (a) H,
 - (b) C₁-C₆ alkyl,
 - (c) aryl, wherein aryl is as defined above,
 - (d) aryl (C₁-C₄) alkyl, where aryl is as defined
 - 15 above,
- or taken together constitute a
- (e) piperidine ring,
 - (f) morpholine ring,
 - (g) piperazine ring, optionally N-substituted with
 - 20 C₁-C₆ alkyl, phenyl or benzyl;

R¹⁷ is

- (a) H,
- (b) C₁-C₆ alkyl,
- (c) phenyl,
- 25 (d) benzyl;

R¹⁸ is

- (a) H,
- (b) C₁-C₆ alkyl,
- (c) phenyl,
- 30 (d) benzyl;

R¹⁹ is

- (a) H,

(b) C₁-C₅ alkyl optionally substituted with a substituent selected from the group consisting of aryl, where aryl is as defined above, heteroaryl where heteroaryl is as defined above, -OH, -SH, C₁-C₄ alkyl, 5 C₁-C₄ alkoxy, C₁-C₄ alkylthio, -CF₃, halo, -NO₂, -CO₂H, -CO₂CH₃, -CO₂-benzyl, -NH₂, C₁-C₄ alkylamino, C₁-C₄ dialkylamino, -PO₃H₂,

(c) aryl, where aryl is defined as above,

(d) -CH₂ aryl, where aryl is defined as above,

10 (e) heteroaryl,

wherein heteroaryl is an unsubstituted, monosubstituted or disubstituted 5- or 6-membered aromatic ring which can optionally contain from 1 to 3 heteroatoms selected from the group consisting of O, N, 15 and S and wherein the substituents are members selected from the group consisting of -OH, -SH, C₁-C₄ alkyl, C₁-C₄ alkoxy, -CF₃, halo, -NO₂, -CO₂H, -CO₂CH₃, -CO₂-benzyl, -NH₂, C₁-C₄ alkylamino, C₁-C₄ dialkylamino; R²⁰ is

20 (a) aryl, where aryl is defined as above,

(b) C₃-C₇ cycloalkyl,

(c) C₁-C₄ perfluoroalkyl,

(d) C₁-C₄ alkyl optionally substituted with a substituent selected from the group consisting of aryl, 25 where aryl is as defined above, heteroaryl where heteroaryl is as defined above, -OH, -SH, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ alkylthio, -CF₃, halo, -NO₂, -CO₂H, -CO₂CH₃, -CO₂-benzyl, -NH₂, C₁-C₄ alkylamino, C₁-C₄ dialkylamino, -PO₃H₂,

30 (e) heteroaryl, where heteroaryl is as defined above;

R²¹ is

(a) H,

(b) C₁-C₆ alkyl,

62

(c) phenyl,

(d) benzyl,

or taken together constitute a

(e) piperidine ring,

5 (f) morpholine ring,

(g) piperazine ring, optionally N-substituted with
C₁-C₆ alkyl, phenyl or benzyl;

X is

10 (a) S,

(b) O,

(c) SO,

(d) SO₂,(e) CHR¹⁴,15 (f) NR¹⁴;

Z is

(a) -O-,

(b) -S-,

20 (c) -NR¹¹-;

m is 1 to 5;

n is 1 to 4;

q is 1 to 5;


t is 2 to 5;

25


or a pharmaceutically acceptable salt thereof.

2. A compound of Claim 1 wherein

30 R¹ is in the para position and is



R^1 is in the para position and is



5

R^6 is


- (a) C_1-C_{10} alkyl,
- (b) C_3-C_{10} alkenyl,
- (c) C_3-C_{10} alkynyl,
- 10 (d) C_3-C_8 cycloalkyl,
- (e) phenyl, optionally substituted with 1-2 substituents selected from the group of halo, C_1-C_4 alkyl, C_1-C_4 alkoxy, $-NO_2$, $-NH_2$, $-OH$ and benzyloxy,
- (f) benzyl, optionally substituted on the phenyl
- 15 ring with one or two substituents selected from the group consisting of halo, C_1-C_4 alkyl, C_1-C_4 alkoxy and $-NO_2$;

R^7 , R^8 , R^9 , R^{10} are independently

- (a) H,
- 20 (b) C_1-C_4 alkyl,
- (c) C_1-C_4 perfluoroalkyl,
- (d) C_3-C_6 cycloalkyl,
- (e) phenyl, optionally substituted with one or two substituents selected from the group of halo, C_1-C_3
- 25 alkyl, C_1-C_3 alkoxy, NO_2 , CF_3 , NH_2 , and OH ,
- (f) R^7 and R^8 taken together are $-(CH_2)_t-$, or $-(CH_2)_mX(CH_2)_q-$,
- (g) R^9 and R^{10} taken together can be S, O;

R¹³ is

- (a) -CONHSO₂R²⁰,
- (b) -NHCONHSO₂R²⁰,
- (c) -NHSO₂NHCOR¹⁹,
- 5 (d) -PO₃H₂,
- (e) -SO₃H,
- (f) -SO₂NHR¹⁹,
- (g) -SO₂NHCOR¹⁹,
- (h) -SO₂NHCONHR¹⁹,
- 10 (i)



or a pharmaceutically acceptable salt thereof.

3. A compound of Claim 2 wherein

15

R⁶ is


- (a) C₁-C₇ alkyl,
 - (b) C₃-C₄ alkenyl,
 - (c) C₃-C₄ alkynyl;
 - 20 (d) phenyl, optionally substituted with 1-2
- substituents selected from the group of halo, C₁-C₄ alkyl, C₁-C₄ alkoxy, -NO₂, -NH₂, -OH and benzyloxy;

R¹³ is

- 25 (a) -CONHSO₂R²⁰,
- (b) -NHCONHSO₂R²⁰,
- (c) -SO₂NHR¹⁹,
- (d) -SO₂NHCONHR¹⁹,
- (e) -SO₂NHCOR¹⁹,
- 30 (f) -SO₂NHCONHR¹⁹;

or a pharmaceutically acceptable salt thereof.

4. A compound of Claim 3 wherein
R¹ is



- 5 or a pharmaceutically acceptable salt thereof.

5. A compound of Claim 4 selected from the group consisting of

- 10 • N-((4'-((4-oxo-2-propyl-8-thia-1,3-diazaspiro-
((4.5))dec-1-en-3-yl-methyl))-((1,1'-biphenyl))-2-
ylsulfonyl))-benzamide
- 15 • N-((4'-((4-oxo-2-propyl-1,3-diazaspiro((4.4))non-
1-en-3-yl)methyl))((1,1'-biphenyl-2-ylsulfonyl))-
benzamide
- 20 • N-((4'-((4-oxo-2-butyl-1,3-diazaspiro((4.4))non-1-
en-3-yl)methyl))((1,1'-biphenyl-2-ylsulfonyl))-
benzamide
- 25 • N-((4'-(4,5-dihydro-4,4-dimethyl-5-oxo-2-propyl-1H-
imidazol-1-ylmethyl))((1,1'-biphenyl))-2-
ylsulfonyl))-benzamide
- N-((4'-(4,5-dihydro-4,4-dimethyl-5-oxo-2-propyl-1H-
imidazol-1-ylmethyl))((1,1'-biphenyl))-2-
ylsulfonyl))-4-chlorobenzamide
- 30 • N-((4'-((4-oxo-2-propyl-8-thia-1,3-diazaspiro-
((4.5))dec-1-en-3y-1-methyl))-((1,1'-biphenyl))-2-
ylsulfonyl))-hexanamide

- N-((4'-((4-oxo-2-propyl-1,3-diazaspiro((2.4))hept-1-en-3-yl)methyl))((1,1'-biphenyl-2-ylsulfonyl))-hexanamide
- 5 • N-((4'-((4-oxo-2-propyl-1,3-diazaspiro((4.4))non-1-en-3-yl)methyl))((1,1'-biphenyl-2-ylsulfonyl))-hexanamide
- 10 • N-((4'-((4-oxo-2-propyl-1,3-diazaspiro((4.5))dec-1-en-3-yl)methyl))((1,1'-biphenyl-2-ylsulfonyl))-hexanamide
- 15 • N-((4'-((4-oxo-2-butyl-1,3-diazaspiro((4.4))non-1-en-3-yl)methyl))((1,1'-biphenyl-2-ylsulfonyl))-hexanamide
- 20 • N-((4'-(4,5-dihydro-4,4-dimethyl-5-oxo-2-propyl-1H-imidazol-1-ylmethyl))((1,1'-biphenyl))-2-ylsulfonyl))-hexanamide
- 25 • N-((4'-(4,5-dihydro-4,4-dicyclopropyl-5-oxo-2-propyl-1H-imidazol-1-ylmethyl))((1,1'-biphenyl))-2-ylsulfonyl))-hexanamide
- 30 • N-((4'-(4,5-dihydro-4,4-bistrifluoromethyl-5-oxo-2-propyl-1H-imidazol-1-ylmethyl))-3'-methyl((1,1'-biphenyl))-2-ylsulfonyl))-hexanamide

- N-((4'-(4,5-dihydro-4,4-dimethyl-5-oxo-2-propyl-1H-imidazol-1-ylmethyl)-4-propyl((1,1'-biphenyl))-2-ylsulfonyl))-hexanamide
- 5 • N-((4'-(4,5-dihydro-4,4-dimethyl-5-oxo-2-propyl-1H-imidazol-1-ylmethyl)((1,1'-biphenyl))-2-ylsulfonyl))-trifluoroacetamide
- 10 • 3,5-dihydro-5,5-dimethyl-2-propyl-3-[(2'-(N-((phenylsulfonyl)carboxamido)biphen-4-yl)methyl]-4H-imidazol-4-one.

6. A pharmaceutical composition comprising a pharmaceutically suitable carrier and a compound of any
15 one of Claims 1 through 4.

7. A pharmaceutical composition comprising a pharmaceutically suitable carrier and a compound of
Claim 5.

20

8. A method of treating hypertension in a warm blooded animal comprising administering to an animal in need of such treatment an effective amount of a compound of any of Claims 1 through 4.

25

9. A method of treating hypertension in a warm blooded animal comprising administering to an animal in need of such treatment an effective amount of a compound of Claim 5.

30

10. A method of treating congestive heart failure in a warm blooded animal comprising administering to an animal in need of such treatment an effective amount of a compound of any of Claims 1 through 4.

11. A method of treating congestive heart failure
in a warm blooded animal comprising administering to an
animal in need of such treatment effective of a compound
5 of Claim 5.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 92/07022

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5	C07D233/70; C07D403/10;	A61K31/415; C07D471/10;
		C07D233/84; C07D491/10;
		C07D235/02 C07D495/10
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System		Classification Symbols
Int.Cl. 5	C07D ;	A61K ; C07F
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A	EP,A,0 380 959 (E.I.DU PONT DE NEMOURS AND COMPANY) 8 August 1990 ---	
A	DE,A,3 913 757 (SCHERING AG) 25 October 1990 ---	
A	EP,A,0 419 048 (MERCK & CO. INC.) 27 March 1991 cited in the application ---	
A	EP,A,0 412 594 (MERCK & CO. INC.) 13 February 1991 cited in the application ---	
A	EP,A,0 291 969 (E.I.DU PONT DE NEMOURS AND COMPANY) 23 November 1988 cited in the application ---	
	-/--	
<p>¹⁰ Special categories of cited documents :</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search 02 DECEMBER 1992		Date of Mailing of this International Search Report 15. 01. 93
International Searching Authority EUROPEAN PATENT OFFICE		Signature of Authorized Officer DE BUYSER I.A.F.

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category °	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
P,X	WO,A,9 114 679 (SANOFI) 3 October 1991 cited in the application see the whole document ---	1,4-11
E	EP,A,0 505 098 (MERCK & CO. INC.) 23 September 1992 see page 3, line 24 see page 5, line 34 see page 5, line 37 -----	1-4,6-11

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

US 9207022
SA 63823

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 02/12/92

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0380959	08-08-90	US-A- 4916129 AU-B- 618503 AU-A- 4854590 CA-A- 2006604 JP-A- 3002169	10-04-90 19-12-91 26-07-90 19-07-90 08-01-91

DE-A-3913757	25-10-90	None	

EP-A-0419048	27-03-91	US-A- 5100897 CA-A- 2024113 JP-A- 3197466	31-03-92 01-03-91 28-08-91

EP-A-0412594	13-02-91	CA-A- 2021954 JP-A- 3148266	29-01-91 25-06-91

EP-A-0291969	23-11-88	US-A- 4820843 AU-B- 603525 AU-A- 1650388 JP-A- 1117876 US-A- 4870186 US-A- 4874867	11-04-89 15-11-90 24-11-88 10-05-89 26-09-89 17-10-89

WO-A-9114679	03-10-91	FR-A- 2659967 FR-A- 2665702 AU-A- 7561091 CA-A- 2057913 EP-A- 0454511	27-09-91 14-02-92 21-10-91 21-09-91 30-10-91

EP-A-0505098	23-09-92	None	
